



DIXIE-DUNDAS FLOOD MITIGATION NATURAL HERITAGE STUDY

Prepared for:

CITY OF MISSISSAUGA

Prepared by:

MATRIX SOLUTIONS INC., A MONTROSE ENVIRONMENTAL COMPANY

Version 2.0

May 2024

Guelph, Ontario

Unit 7B, 650 Woodlawn Road West

Guelph, Ontario, Canada N1K 1B8

T 519.772.3777 F 226.314.1908

www.matrix-solutions.com

**DIXIE-DUNDAS FLOOD MITIGATION
NATURAL HERITAGE STUDY**

Prepared for City of Mississauga, May 2024

Andrea Perez, B.Sc.
Ecologist

Shari Muscat, B.A., B.E.S.
Senior Environmental Planner

CONTRIBUTORS

Name	Job Title	Role
Amanda McKay, P.Eng., PMP	Water Resources Engineer	Co-author
Andrea Perez, B.Sc.	Ecologist	Co-author
Shari Muscat, B.A., B.E.S.	Senior Environmental Planner	Reviewer
J. Arnel Fausto, M.Sc.	Senior Ecologist	Reviewer

DISCLAIMER

Matrix Solutions Inc. certifies that this report is accurate and complete and accords with the information available during the project. Information obtained during the project or provided by third parties is believed to be accurate but is not guaranteed. Matrix Solutions Inc. has exercised reasonable skill, care, and diligence in assessing the information obtained during the preparation of this report.

This report was prepared for the City of Mississauga. The report may not be relied upon by any other person or entity without the written consent of Matrix Solutions Inc. and of the City of Mississauga. Any uses of this report by a third party or any reliance on decisions made based on it are the responsibility of that party. Neither Matrix Solutions Inc. nor its affiliates are responsible for damages or injuries incurred by any third party as a result of decisions made or actions taken based on this report.

VERSION CONTROL

Version	Date	Issue Type	Filename	Description
V0.1	19-Mar-2020	Draft	24604-531 R NHS 2020-03-19 draft V0.1.docx	Issued to client for review
V1.0	08-Jul-2020	Final	24604-531 R NHS 2020-07-08 final V1.0.docx	Issued to client
V1.1	30-Nov-2021	Draft revised	24604-531 R NHS 2020-11-30 draft V1.1.docx	Report updated to incorporate study area south of Dundas Street; revisions issued to client for review
V1.2	21-Nov-2023	Draft revised 1	24604-531 R NHS 2023-11-21 draft V1.2.docx	Updates included from client; reissued for review
V1.3	15-Mar-2024	Draft revised 2	24604-531 R NHS 2024-03-15 draft V1.3.docx	Addressed agency comments. Updated with preferred alternatives.
V2.0	29-May-2024	Revised Final	24604-531 R NHS 2024-05-29 final V2.0.docx	Minor update to reflect changes to <i>Conservation Authorities Act</i> . Issued to client.

TABLE OF CONTENTS

1	INTRODUCTION.....	1
1.1	Study Area.....	1
2	REGULATORY FRAMEWORK.....	3
3	BACKGROUND REVIEW	9
3.1	Methods.....	9
3.1.1	Agency Correspondence	9
3.1.2	Background Sources.....	9
3.2	Analysis of Significance and Sensitivity.....	10
3.2.1	Natural Area Designations	10
3.2.2	Species at Risk Screening	10
4	FIELD METHODOLOGY	11
4.1.1	Ecological Land Classification.....	12
4.1.2	Botanical Inventories	12
4.1.3	Bat Maternity Roosting Survey	12
4.1.4	Incidental Wildlife	13
4.1.5	Species at Risk Assessment.....	13
4.1.6	Fish Habitat Assessment	14
4.1.7	Fish Community Assessment	15
5	EXISTING NATURAL ENVIRONMENT	17
5.1	Topography.....	17
5.2	Terrestrial Habitat.....	17
5.2.1	Vegetation Communities	17
5.2.2	Flora	21
5.2.3	Incidental Wildlife Observations.....	22
5.3	Aquatic Resources.....	23
5.3.1	Aquatic Habitat	23
5.3.2	Fish Community	32
6	SIGNIFICANT NATURAL HERITAGE FEATURES AND FUNCTIONS	33
6.1	Significant Valleylands and Corridors	33
6.2	Significant Woodlands	34
6.3	Fish and Fish Habitat.....	34
6.4	Linkages and Corridors.....	34
6.5	Species at Risk.....	34
6.5.1	Confirmed SAR	35
6.5.2	Moderate Probability SAR	35
6.5.3	Confirmed SCC	36
6.5.4	Moderate Probability SCC.....	36
6.5.5	Bat Maternity Roosting Survey	36
6.6	Significant Features and Functions Summary.....	40

7	DESCRIPTION OF ALTERNATIVES	42
7.1	Dixie Area Alternative Solutions	42
7.1.1	Dixie AS1: Improved Conveyance with a Minimized Footprint	42
7.1.2	Dixie AS2: Improved Conveyance by Making Room for Little Etobicoke Creek ..	42
7.1.3	Dixie AS3: Flood Containment with Mitigation for Upstream Impacts	42
7.1.4	Dixie Do Nothing	43
7.1.5	Preferred Alternative	43
7.2	Dundas Area Alternative Solutions	43
7.2.1	Dundas AS1	43
7.2.2	Dundas AS2	44
7.2.3	Dundas AS3	44
7.2.4	Dundas Do Nothing.....	44
7.2.5	Preferred Alternative	45
8	RECOMMENDATIONS AND PERMITTING	45
9	CONCLUSION.....	46
10	REFERENCES	46

IN-TEXT FIGURES

Figure 1	Study Area.....	2
Figure 2	Fish Sampling Locations	16
Figure 3	Ecological Land Classification.....	20
Figure 4a	Fish Habitat Upstream of Dixie Road	25
Figure 4b	Fish Habitat Downstream of Dixie Road	27
Figure 4c	Fish Habitat Upstream of Dundas Street East	29
Figure 4d	Fish Habitat downstream of Dundas Street East	31
Figure 5	Bat Maternity Roosting Snags within the Study Area.....	39
Figure 6	Natural Heritage Features and Buffers	41

IN-TEXT TABLES

Table 1	Applicable Federal, Provincial, and Municipal Policies.....	4
Table 2	Background Sources.....	9
Table 3	Field Survey Summary.....	11
Table 4	Ecological Land Classification Communities	17
Table 5	Floristic Summary and Assessment of Results.....	22
Table 6	Historical Fisheries Data at Little Etobicoke Creek Within and Surrounding the Study Area.....	32
Table 7	Fish Community Results for Little Etobicoke Creek in 2019 and 2021	33
Table 8	Summary of Little Brown Myotis and Northern Myotis Suitable Roost Trees within Study Area (Leaf-off Survey)	37
Table 9	Summary of Tri-Coloured Bat Suitable Roost Trees within Study Area (Leaf-on Survey)	38

APPENDICES

APPENDIX A	Wildlife Lists
APPENDIX B	Site Photographs
APPENDIX C	Species at Risk and Species of Conservation Concern Screening

1 INTRODUCTION


Matrix Solutions Inc., a Montrose Environmental company, was retained by the City of Mississauga (the “City”) to create a feasibility study for flood mitigation measures in Little Etobicoke Creek. Little Etobicoke Creek is an urban creek system which has been highly impacted by flood flows in recent years. Under regulatory flow conditions, approximately 130 m³/s of flood water spills beyond the floodplain of Little Etobicoke Creek at Queen Frederica Drive. And as a result, over 1,000 residential and commercial buildings between the Queen Frederica Drive and the Queen Elizabeth Way (QEW) are at risk of flood damage.










The City’s objective for the flood mitigation feasibility study is to modify the natural and built infrastructure of Little Etobicoke Creek, such that flood flows are contained within the floodplain upstream of Dixie Road, thereby removing the spill that flows toward the Applewood Creek watershed during large rainfall events. To assist the project through the various phases of the Municipal Class Environmental Assessment process, various technical studies were completed. This Natural Heritage Study (NHS) Report is one of those technical studies and has identified the existing conditions of the natural heritage features within the study area. The NHS report consists of data collected during background reviews and field studies, as well as an assessment of the environmental impacts of the alternative solutions.

1.1 Study Area

The study area is located within Little Etobicoke Creek between the area 500 m west of Dixie Road and approximately 360 m east of Dundas Street East within the City (Figure 1). Originally, the project study area encompassed only the area approximately 500 m west of Dixie Road and 200 m east of Dixie Road; however, two study area extensions have occurred. First, in the summer of 2019 the study area was extended to include the entire area of Little Etobicoke Creek all the way to Dundas Street East. The second extension occurred in 2021 with the addition of the 360 m of creek east of Dundas Street East.

Little Etobicoke Creek is part of the Etobicoke Creek watershed which is regulated by the Toronto and Region Conservation Authority (TRCA) and is also part of the City’s “Significant Natural Areas and Natural Green Spaces.” The large meander bend at the northeast corner of the site, approximately 500 m upstream of Dundas Street East, is considered a “Special Management Area” within the City (Figure 1). “Special Management Areas are lands adjacent to or near Significant Natural Areas or Natural Green Spaces and will be managed or restored to enhance and support the Significant Natural Area or Natural Green Space” (City of Mississauga 2021). Within the study area, Little Etobicoke Creek is surrounded by a small, predominantly deciduous, riparian community as well as residential and commercial buildings.



-  Natural Heritage Study Area
-  Special Policy Area
-  Special Management Areas
-  Significant Natural Areas and Natural Green Spaces
-  Watercourse
-  Flow Direction
-  Highway
-  Road
-  Dixie Road Bridge



City of Mississauga
Dixie-Dundas Flood Mitigation

Natural Heritage Study Area

Date: November 2023	Project: 24603	Submitter: E. Wilkinson	Reviewer: R. Leppington
---------------------	----------------	-------------------------	-------------------------

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.

I:\CityOfMississauga\24603\FiguresandTables\ECL2019\Report\Figure-1_Study_Area.mxd - Tabloid_L - 16-Nov-23, 10:12 AM - s.yam - TID005

2 REGULATORY FRAMEWORK

Review of the regulatory framework provides guidance on the protection of natural heritage features and evaluation of significance. Natural heritage features identified within the study area were evaluated against the federal, provincial, and municipal planning policies applicable to the study area (Table 1).

Table 1 Applicable Federal, Provincial, and Municipal Policies

Legislation	Policies and Regulations	Guidelines	Summary of Contents
Federal Acts and Regulations			
<i>Species at Risk Act</i> (SARA; 2002)	N/A	N/A	<ul style="list-style-type: none"> SARA incorporates a number of prohibitions to protect SAR, providing recovery strategies for extirpated, endangered, or threatened species, as well as managing species of special concern (Government of Canada, 2016). On private or provincially-owned lands, only aquatic species listed as endangered, threatened, or extirpated and migratory birds listed on Schedule 1 are protected under SARA, unless ordered by the Governor in Council, or for those species that have critical habitat identified. Schedule 1 migratory birds where critical habitat has been identified require consideration under SARA. Should a species also be listed under SARA and the ESA, where the ESA provides equal or greater protection, the ESA takes precedence. Applicability to Project: Should any SAR fish, mussel, or migratory bird species be identified within the study area, protections under SARA will come into effect.
<i>Fisheries Act</i> (1985, amended 2019)	Fish Protection Policy Statement (2013)	N/A	<ul style="list-style-type: none"> The <i>Fisheries Act</i> outlines the framework for the management and regulation of fisheries and the conservation and protection of fish and fish habitat within the fishing zones of Canada, all waters in the territorial sea of Canada, and all internal waters of Canada. The most recent revision (2019) restores protections to both fish and fish habitat by prohibiting the harmful alteration, disruption, or destruction (HADD) of fish habitat and prohibiting the death of fish. Environmental recommendations and mitigation measures are to include the provision of timing windows, to avoid the death of fish or the HADD of fish habitat. Any proposed works that may result in either require a Request for Review submitted to the DFO. Applicability to Project: Any activities that may impact Little Etobicoke Creek or other watercourses within the study area could require a Request

Legislation	Policies and Regulations	Guidelines	Summary of Contents
			for Review from the DFO. In-water work is not permitted within the watercourse during the fisheries timing window for warmwater fish from April 1 to June 30.
<i>Migratory Birds Convention Act</i> (MBCA; 1994, amended on 2017)	Migratory Birds Regulation (2022) Migratory Bird Sanctuary Regulations (2022)	N/A	<ul style="list-style-type: none"> The MBCA affords protection to birds listed under Article 1 of the MBCA. General prohibitions protect migratory birds, their nests, and their eggs, and prohibits the deposition of harmful substances in waters and other areas frequented by them. The MBR clarifies the nest protection timelines for 18 species that are listed on Schedule 1. These timelines include a waiting period until the nest can be deemed abandoned and subsequently removed. The waiting period for these 18 species remains in effect even if the nest is unoccupied. Nest protection for all other species which are not listed in Article 1 remains the same, with nest removal occurring once the young have fledged. Applicability to Project: All vegetation removal shall occur outside of the migratory bird nesting window of April 1 to August 31 of any year to avoid disturbing active nests.
Provincial Acts and Regulations			
<i>Places to Grow Act</i> (2005)	<i>A Place to Grow. Growth Plan for the Greater Golden Horseshoe</i> (2020) O. Reg. 416/05: Growth Plan Areas	N/A	<ul style="list-style-type: none"> The <i>Places to Grow Act</i> enables the development of regional growth plans that guide government investments and land use planning policies. The Growth Plan for the Greater Golden Horseshoe provides a framework for long-term management of growth within the region. It builds upon the policies in the PPS 2020 provides direction on how and where the region will grow. Applicability to Project: The Region of Peel has been designated as a Growth Plan Area. The project area is subject to policies under the Growth Plan for the Greater Golden Horseshoe (2020).

Legislation	Policies and Regulations	Guidelines	Summary of Contents
<i>Endangered Species Act (ESA; 2007, amended 2020)</i>	O. Reg.'s: 230/08 242/08 829/21 830/21 832/21	N/A	<ul style="list-style-type: none"> The ESA provides conservation and protection of species in Ontario. Species listed as endangered or threatened under the ESA are afforded legal protection from harm and harassment. The ESA also prohibits damage or destruction of habitat of endangered or threatened species. Habitat protection for a species can be general or subject to the specific provisions of a habitat regulation as set out in O. Reg. 832/21. Applicability to Project: The ESA applies to all SAR within provincial lands protected under the ESA. Any impacts to these species or their habitats protected under the ESA would require a permit. There is a confirmed Butternut within the study area and there is potential for other SAR to occur within it as well.
<i>Fish and Wildlife Conservation Act (1997, amended 2021)</i>	O. Reg.'s: 663/98 664/98 665/98 666/98 667/98 668/98 669/98 670/98	N/A	<ul style="list-style-type: none"> The <i>Fish and Wildlife Conservation Act</i> affords protection for some species of birds, amphibians, reptiles, and mammals in Ontario. Some bird species which are not afforded protection under the MBCA are afforded protection under the <i>Fish and Wildlife Conservation Act</i>. Applicability to Project: There may be suitable habitat along Little Etobicoke Creek, and within the study area for some of these species, such as raptors or fur bearing mammals. Environmental management considerations should include timing windows for vegetation clearing and for construction.
<i>Conservation Authorities Act (1990, amended 2022)</i>	O. Reg. 41/24: Prohibited Activities, Exemptions and Permits The Living City Policies for Planning and Development in	Planning & Development Procedural Manual (TRCA, 2008) TRCA Environmental Impact Statement Guidelines (2014)	<ul style="list-style-type: none"> The <i>Conservation Authorities Act</i> empowers conservation authorities to regulate development within hazardous lands, wetlands, river or stream valleys; within areas that are adjacent to the shoreline of the Great Lakes; inland lakes subject to flooding, erosion, or dynamic beach hazards; as well as any activities that interfere with an existing watercourse or wetland. On April 1, 2024, portions of the <i>Conservation Authorities Act</i> were repealed and substituted, including but not limited to Part VI (Section 28 and Section 29) and Part VII. On April 1, 2024, O. Reg. 41/24 was enacted under the <i>Conservation</i>

Legislation	Policies and Regulations	Guidelines	Summary of Contents
	the Watersheds of the Toronto and Region Conservation Authority (TRCA, 2014)	<p>Guideline for Determining Ecosystem Compensation (TRCA, 2018)</p> <p>Erosion and Sediment Control Guide for Urban Construction (TRCA, 2019)</p> <p>Seed Mix Guideline V.2.0 (TRCA, 2022)</p>	<p><i>Authorities Act</i>, which supersedes previous Conservation Authority specific Regulation's of Development, such as TRCA's O. Reg 166/06. O. Reg. 41/24 includes updated definitions for key terms; an updated list of prohibited activities, or activities that would require permission; clarification on the pre-consultation requirements for permits; and updates to floodplain and regulated area mapping standards. Notably, the regulated areas surrounding wetlands has been updated to a standard 30 m regulation limit for all wetlands (including PSW's and wetlands greater than 2ha in size).</p> <p>Applicability to Project:</p> <ul style="list-style-type: none"> • A portion of the study area is located within the jurisdiction of the TRCA (regulated under O. Reg. 41/24). Any works within the regulatory limit may require a permit. • A portion of the project is located within two Special Policy Areas and is subject to additional provincial regulations. <p>Additional Guidelines:</p> <ul style="list-style-type: none"> • Crossing Guidelines for Valley and Stream Corridors (TRCA, 2015) • Forest Edge Management Plan Guidelines (TRCA, 2004) • Flora Species Native to the TRCA Jurisdiction (TRCA, 2022) • Preserving and Restoring Healthy Soil: Best Practices for Urban Construction (TRCA, 2012)
Municipal Acts and Regulations			
<i>Region of Peel Official Plan (2022)</i>	N/A	N/A	<ul style="list-style-type: none"> • The Greenlands System identifies Core Natural Heritage features and linkages within the landscape. These include wetlands, woodlands, valleylands, stream corridors, shorelines and lakes, ANSIs, significant wildlife habitat and other natural or protected areas. • Applicability to Project: The project area contains core areas as part of the Greenlands System. These lands are subject to the natural heritage policies as outlined in the Region of Peel Official Plan.

Legislation	Policies and Regulations	Guidelines	Summary of Contents
<i>City of Mississauga Official Plan (July 2023 Consolidation)</i>	N/A	N/A	<ul style="list-style-type: none"> Mississauga’s Green System is comprised of the natural heritage system, urban forest, natural hazard lands, and parks and open spaces. The City is committed to protecting, enhancing, restoring, and expanding its Green System. Applicability to Project: The study area is part of the Green System, and is mapped as a Natural Hazard, Special Policy Area Floodplain, Special Management Area, and Significant Natural Area and Natural Green Space.

Notes:

- ANSI - Area of Natural and Scientific Interest
- DFO - Fisheries and Oceans Canada
- ESA - *Endangered Species Act*
- TRCA – Toronto and Region Conservation Authority
- HADD - harmful alteration, destruction, and displacement
- MBCA - *Migratory Birds Convention Act*
- MBR - Migratory Birds Regulation
- NHRM - *Natural Heritage Reference Manual*
- O. Reg. - Ontario Regulation
- PPS - Provincial Policy Statement
- SAR - species at risk
- SARA - *Species at Risk Act*

3 BACKGROUND REVIEW

3.1 Methods

The following information sources were reviewed for records related to natural heritage features that have the potential or are known to occur within the study area.

3.1.1 Agency Correspondence

Initial information requests regarding terrestrial sensitivities and species at risk (SAR) were submitted to the Ministry of Environment, Conservation and Parks (MECP), and to the TRCA. A background request for natural heritage information was submitted to the TRCA on June 18, 2019. Information was received on June 15, 2019, and was incorporated into Appendix A. A project screening request was sent to the MECP on May 15, 2019, for information related to natural heritage features and SAR potential within the study area. The MECP responded on May 15, 2019, indicating that there was no additional information to be provided. An updated screening request was sent to MECP on April 13, 2021, to include the portion of study area downstream of Dundas Street East. The MECP responded on April 22, 2021, and provided a list of 11 SAR potentially within the study area.

3.1.2 Background Sources

In addition to information provided by these regulatory agencies, other publicly available data sources were reviewed to determine potential Species of Conservation Concern (SCC) and SAR whose occurrence ranges overlap with the study area. Background review material for the study area has also been obtained from available secondary source reports. The majority of background information was provided by the TRCA. Table 2 summarizes the complete list of background information reviewed for the study area.

Table 2 Background Sources

Source	Citation	Information Reviewed
Land Information Ontario (LIO)	MNR 2000	Natural heritage features data layers
Ontario GeoHub	MNRF 2024a	Natural heritage features data layers
Aquatic Species at Risk Map	DFO 2024	Aquatic species at risk map
Natural Heritage Information Centre (NHIC)	MNRF 2024b	Data records for the study area
Ontario Breeding Bird Atlas (OBBA)	Birds Canada et al. 2024	Species records for the study area
Ontario Butterfly Atlas	TEA 2024a	Species records for the study area
Ontario Moth Atlas	TEA 2024b	Species records for the study area
Ontario Reptile and Amphibian Atlas	Ontario Nature 2024	Species records for the study area
Important Bird Areas	Bird Studies Canada 2024	Data records for the study area

Source	Citation	Information Reviewed
eBird	Cornell Lab of Ornithology 2024	Species records for the study area
iNaturalist	CAS 2024a	Species records for the study area
Ontario Mammals, iNaturalist	CAS 2024b	Species records for the study area
Etobicoke and Mimico Creeks Watersheds Technical Report	TRCA 2010	Species records and natural heritage features for the study area
Etobicoke Creek Watershed Report Card	TRCA 2018	Overall health of watershed
Etobicoke Creek - The Aquatic Ecosystem	TRCA 2006	Species records and natural heritage features for the study area
Natural Areas Survey (NAS)	City of Mississauga 2020	Natural areas across the city

3.2 Analysis of Significance and Sensitivity

The ecological features identified within the study area were evaluated to determine the significance of each feature. Significance is based on regional, provincial, and federal designations, which are described in the following subsections.

3.2.1 Natural Area Designations

Natural area designations are those that are recognized as significant on official plans or in other policy planning documents. This includes Areas of Natural or Scientific Interest (ANSIs; provincially, regionally, or other), significant wetlands (provincially, regionally, or locally), significant woodlands, and Environmentally Significant Areas. ANSIs and Environmentally Significant Areas are evaluated by the province or municipality. Only wetlands and woodlands can be assessed for significance by non-government organizations.

3.2.2 Species at Risk Screening

The background review identified potential SAR that could occur within the study area. All SAR identified were screened to determine the likelihood of occurrence and whether suitable habitat is present.

SAR are defined in this report to include the following provincial and federal designations:

- ESA (provincial; Government of Ontario 2021a): all provincially designated species that are listed as Extirpated, Endangered, or Threatened on the SARO List and protected under the ESA; species listed as Special Concern are considered a SCC, as they are not protected under the ESA.

- SARA (federal; Government of Canada 2021): only applies to fish and migratory birds protected under the Migratory Birds Convention Act, 1994 (MBCA; Government of Canada 2017), anywhere they occur (e.g., includes non-federal land), that are designated as Extirpated, Endangered, and/or Threatened under the SARA. All other species are only protected if special provisions or executive orders are made.

To determine if suitable habitat for SAR is available within the study area, the preferred habitat requirements for reported SAR were compared to vegetation communities, aquatic habitats, and niche habitats identified during field inventories and the background review. The results of the SAR habitat screening are provided in Section 6.5.

4 FIELD METHODOLOGY

Field inventories were completed within the study area by Matrix staff during the spring and summer of 2019 as part of the NHS. Additional field studies were completed in the spring of 2020 to capture spring botanicals within the extended study area up to Dundas Street East. Additional field studies were completed in the spring, summer, and fall of 2021 to obtain data on the extended study area east of Dundas Street East. The names of staff members and the field inventories that they completed are provided in Table 3.

Table 3 Field Survey Summary

Field Inventory	Date	Matrix Staff
Vegetation (Ecological Land Classification, Botanical Inventory, Invasive Species)	Spring May 24, 2019	K. Reis
	Summer July 10, 2019	K. Reis and B. MacVeigh
	Summer August 22, 2019	K. Reis and E. Wilkinson
	Fall September 25, 2019	E. Wilkinson
	Spring April 8, 2020	E. Wilkinson
	Spring April 16, 2021	E. Wilkinson
	Summer July 30, 2021	E. Wilkinson
	Fall September 24, 2021	E. Wilkinson
Fish and Fish Habitat	July 10, 2019	K. Reis and B. MacVeigh
	August 22, 2019	K. Reis and E. Wilkinson
	July 30, 2021	R. Leppington and K. Reich
Bat Maternity Roosting Habitat Survey	April 8, 2020 (Leaf-off)	K. Reis and E. Wilkinson
	June 16, 2020 (Leaf-on)	E. Wilkinson
	April 16, 2021 (Leaf-off)	E. Wilkinson
	July 30, 2021 (Leaf-on)	E. Wilkinson
Incidental Observations	May 24, 2019	K. Reis
	July 10, 2019	K. Reis and B. MacVeigh
	August 22, 2019	K. Reis and E. Wilkinson
	September 25, 2019	K. Reis and E. Wilkinson
	April 16, 2021	E. Wilkinson
	July 20, 2021	E. Wilkinson, K. Reich and R. Leppington
	September 24, 2021	E. Wilkinson and K. Reis

4.1.1 Ecological Land Classification

Vegetation community delineation was completed within the study area using aerial photography and refined thorough investigations in the field. The standard Ecological Land Classification (ELC) system for southern Ontario (Lee 2008; Lee et al. 1998) as well as the TRCA Ecological Land Classification Codes and Common Names (TRCA 2019a) were applied. Details of the vegetation communities that were recorded included species composition and dominance, community structure, uncommon species or features, and evidence of anthropogenic disturbance. Vegetation community status rarity was assessed through NHIC vegetation community rankings (MNR 2019) and the local rarity rankings in the Annual Local Occurrence Score and Local Rank Update (TRCA 2017).

4.1.2 Botanical Inventories

A botanical inventory was completed during the field inventories for each of the vegetation communities. The field investigations were completed during spring, summer, and fall (Table 3). A list of species was compiled to determine the presence of SCC, SAR, and invasive species. Habitats of SCC, SAR, and invasive species identified during the field inventories were mapped for the ELC community in which they encompassed.

Plants were identified to family, genus, species, subspecies, and hybrid level according to the Newmaster (1998) Ontario Plant List and cross-referenced with the Database of Vascular Plants of Canada (VASCAN; Brouillet et al. 2020) for scientifically accepted nomenclature.

4.1.3 Bat Maternity Roosting Survey

The location of suitable bat maternity roosting habitat, including snags, was identified following the modified methodology of the Guelph District Survey Protocol for Species at Risk Bats within Treed Habitats: Little Brown Myotis, Northern Myotis, and Tri-colored Bat (MNR 2017) and Bats and Bat Habitat: Guidelines for Wind Power Projects (MNR 2011). This scoped assessment will provide the likelihood that appropriate habitat for SAR bats is present, however it will not confirm the presence or absence of any bat species.

Phase 1 Bat Habitat Suitability Assessment consists of evaluating the study area and deciding whether any area would be designated as a coniferous, deciduous, or mixed wooded ELC ecosite. Based on aerial photography the study area is dominated by deciduous trees and is adjacent to a large watercourse, these two factors provide a potential suitable habitat for bat maternity roosts. Phase 2 was therefore initiated and was divided into a leaf-off and a leaf-on assessment. Identifying suitable roost trees for Little Brown Myotis and Northern Myotis includes recording the location of all snags that exhibit appropriate attributes including cavities, loose bark, cracks, or knot holes. Identifying suitable roost trees for Tri-Colored Bats includes recording the location of any Oak trees greater than 10 cm diameter at breast height (DBH), Maple trees greater than 10 cm DBH if the tree includes dead/dying leaf clusters, and any Maple tree greater than 25 cm DBH. Dates of the field inventories are provided in Table 3.

4.1.4 Incidental Wildlife

All wildlife observations were documented on all field visits. This included actual direct observations (including vocalizations) of individuals as well as signs of wildlife presence (i.e., tracks, scats, dens, nests, etc.).

4.1.5 Species at Risk Assessment

A habitat screening was conducted to determine the potential occurrence of SAR within the study area. For this screening, SAR are defined as species that are listed as either THR or END under the ESA. Individuals of these species, as well as their habitat, are protected in Ontario. Species listed as SC under the ESA receive protection under the NHRM and their habitat is considered SWH. Species listed under SARA are only protected on federal land or as part of projects that are otherwise being permitted by a federal agency. This includes aquatic SAR.

Matrix conducted a background records review to create a list of potential SAR species within the study area. This list was developed using publicly available resources and community-based ecological monitoring databases. Once the list was compiled, a screening exercise was completed to determine the presence of suitable habitat for each SAR identified as potentially occurring within the subject lands, based on known preferred habitat characteristics for each species. The following species were included within the SAR Screening:

- Species listed as either Threatened (THR) or Endangered (END) under the Endangered Species Act (ESA) administered by the MECP. Individuals of these species, as well as their habitat, are protected in Ontario.
- SCC include those listed as SC under ESA, or species listed as threatened or endangered under the federal Species at Risk Act (SARA), not protected under the ESA, and species with Provincial S-rank assigned by the NHIC as S1 (critically imperiled), S2 (imperiled) or S3 (vulnerable). SCC receive protection under the Natural Heritage Reference Manual, and their habitat is considered significant wildlife habitat (SWH).
- Species listed under SARA are only protected on federal land, as part of projects that are otherwise being permitted by a federal agency and includes aquatic SAR. SARA also requires consideration for any migratory bird listed on Schedule 1 where critical habitat has been identified; however, should the species also be listed under ESA and provides equal or greater protection, the ESA take precedence.

The following probability rankings were used to assess potential habitat within the study area:

- **Low Probability:** The site lacks the necessary size, geographic location, or other features required for SAR habitat.

- **Moderate Probability:** The site contains candidate habitat features, as identified during the site visit and through aerial photography interpretation. Species records identified the species within 10 km of the study area.
- **Confirmed:** The site contains candidate habitat features, as identified during the site visit and through aerial photography interpretation and the SAR was observed onsite by field staff.

4.1.6 Fish Habitat Assessment

A qualitative assessment of the habitat potential based on a modified Ontario Stream Assessment Protocol (OSAP; Stanfield 2017) was conducted in Little Etobicoke Creek within the study area. The objective of this assessment was to characterize the local aquatic habitat and assign a qualitative habitat potential ranking. Characteristics of high-quality aquatic habitat include natural sinuosity with a well-defined riffle/pool sequence, variability in water depth and bed substrate, naturally occurring woody debris, undercut banks, and natural riparian vegetation overhanging the banks that provides food for various aquatic organisms. The greater the quantity of preferred habitat features present, the higher potential aquatic habitat ranking. The creek was inventoried throughout the reach for a variety of geomorphic features (i.e., riffles, pools, and runs). The modified qualitative OSAP approach included documentation and assessment of the following watercourse conditions:


- general watercourse characteristics (i.e., stream pattern, general gradient, and flow)
- channel characteristics (i.e., wetted width and depth, bankfull width and depth, and depth of riffles/pools/run)
- substrate and bank materials
- other pertinent habitat features (i.e., spawning, nursery, and refuge areas, barriers to fish movement, and macrophyte growth)
- disturbances and evidence of past habitat alterations (i.e., channelization, channel hardening, or straightening)

After the completion of the aquatic habitat assessment, field data were summarized to determine the overall habitat potential.

4.1.7 Fish Community Assessment

Fish community sampling was conducted at five sampling stations (Figure 2). Prior to sampling, a Licence to Collect Fish for Scientific Purposes was obtained from the Aurora District MNR on June 3, 2019, and was amended on June 9, 2019, to include the study area up to Dundas Street East. A follow up licence was obtained on April 23, 2021, to complete fish community sampling in the study area east of Dundas Street East. Following the OSAP Section 3 Module 1 (Stanfield 2017), a single-pass backpack electrofishing survey was conducted, sampling all available habitat within the sampling station. As per OSAP, sampling stations were a minimum of 40 m in length and contained at least one riffle/pool sequence. The amount of effort expended at each sampling station was dependent on the total area of the station.

After fish were captured, fish were transferred to holding bins and processed. All fish with well-defined identification markings or morphometric characteristics were enumerated (i.e., identified, measured, and weighed) and released back to the creek. Electrofisher shocker seconds and shock times were also recorded.



- Natural Heritage Study Area
- Watercourse
- Flow Direction
- Highway
- Road
- Dixie Road Bridge
- Fish Sampling Site

Reference: Contains information licensed under the Open Government Licence – Ontario. Contains data obtained by Matrix Solutions Inc. and the City of Mississauga. Imagery: Service Layer Credits: © 2023 Microsoft Corporation © 2023 Maxar © CNES (2023) Distribution Airbus DS

1:5,000 metres
50 0 50 100
NAD 1983 UTM Zone 17N



City of Mississauga
Dixie-Dundas Flood Mitigation

Fish Sampling Sites

Date: November 2023	Project: 24603	Submitter: E. Wilkinson	Reviewer: R. Leppington
---------------------	----------------	-------------------------	-------------------------

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.

I:\CityOfMississauga\24603\FiguresandTables\ECL2019\Report\Figure-2\Fish_Sampling_Locations.mxd - Tabloid_L - 16-Nov-23, 10:13 AM - slyam - T10005

5 EXISTING NATURAL ENVIRONMENT

Characterization of the natural environment is provided in the following subsections. A complete list of species observed during field investigations, as well as in the background review, is located in Appendix A. Site photographs are presented in Appendix B.

5.1 Topography

The major physiographic regions in the Etobicoke watershed include the South Slope and Lake Iroquois Sand Plain. The South Slope begins south of the Oak Ridges Moraine and slopes toward Lake Ontario. It is characterized by a smooth, faintly drumlinized, clay till plain, and a deeply incised stream valley (TRCA 2010). The Lake Iroquois Sand Plain is comprised of sand, silt, and clay deposits, with the finer materials being closer to the current Lake Ontario shoreline (TRCA 2010). The geology of Little Etobicoke Creek is comprised predominately of clay loam within the upper reaches of the creek and sandy loam, sand, and clay near the mouth of the creek (TRCA 2010).

The watershed consists of three major land uses, including: 63% urban, 22% rural, and 15% natural land cover. The majority of urban land use is made up of industrial, commercial, and residential buildings occupying approximately 49% of the watershed (TRCA 2010).

5.2 Terrestrial Habitat

5.2.1 Vegetation Communities


Vegetation communities within the study area are mapped on Figure 3 and described in further detail in Table 4. ELC data collected through the TRCA was available for the study area, with the exception of the study area east of Dundas Street East. The TRCA ELC data was confirmed, where available, and ELC data was collected in the area east of Dundas Street East to determine vegetation communities. In total, eight terrestrial ELC communities and one aquatic community were documented.

Table 4 Ecological Land Classification Communities

Ecological Land Classification Community Type	Community Description
FOD7-a Fresh-Moist Manitoba Maple Lowland Deciduous Forest Local Rank: L5	This community was characterized by a dominant Manitoba Maple (<i>Acer negundo</i>) canopy and subcanopy followed by Norway Maple (<i>Acer platanoides</i>) and Common Buckthorn (<i>Rhamnus cathartica</i>). The ground cover consisted primarily of non-native species such as Garlic Mustard (<i>Alliaria petiolate</i>) and Sweet Violet (<i>Viola odorata</i>) as well as native Goldenrod species (<i>Solidago sp.</i>). This community has a local rank of L5 indicating this community is common and secure within the region.
FOD7-b	This community was characterized by Norway Maple and Manitoba Maple within the canopy and subcanopy followed by

Ecological Land Classification Community Type	Community Description
Fresh-Moist Norway Maple Lowland Deciduous Forest Local Rank: L+	Goldenrod species and Garlic Mustard within the groundcover layer. This community has a local rank of L+ indicating this community is dominated by non-native species within the canopy.
FOD7-3 Fresh-Moist Willow Lowland Deciduous Forest Local Rank: L5	This community was characterized by Willow species (<i>Salix sp.</i>) and Manitoba Maple within the canopy followed by Manitoba Maple, Norway Maple and Green Ash (<i>Fraxinus pennsylvanica</i>) within the subcanopy. The understory and ground cover were comprised of Virginia Creeper (<i>Parthenocissus quinquefolia</i>), Bittersweet Nightshade (<i>Solanum dulcamara</i>), Red Raspberry (<i>Rubus idaeus</i>), and Goldenrod species. This community has a local rank of L5 indicating this community is common and secure within the region.
FOD7-2 Fresh-moist Ash Deciduous Forest Local Rank: L4	This community was characterized by Green Ash and Manitoba Maple within the canopy and Norway Maple, and White Mulberry (<i>Morus alba</i>) within the subcanopy. The understory and groundcover consisted of Virginia Creeper, Green Ash saplings, and Goldenrod species. This community has a local rank of L4 indicating that this community is generally secure in the rural matrix, but of conservation concern in the urban matrix.
CUP2-A Restoration Mixed Plantation Local Rank: L5	This community was characterized by Red Pine (<i>Pinus resinosa</i>), Manitoba Maple and Green Ash within the canopy and subcanopy. The understory and groundcover consisted of Wild Carrot (<i>Daucus carota</i>), Cow Vetch (<i>Vicia cracca</i>), and Virginia Creeper. This community has a local rank of L5 indicating this community is common and secure within the region.
CUW1-A3 Native Deciduous Successional Woodland Local Rank: L5	This community was characterized by Red Oak (<i>Quercus rubra</i>) and White Birch (<i>Betula papyrifera</i>) within the sparse canopy and Basswood (<i>Tilia Americana</i>), and Blue Beech (<i>Carpinus caroliniana</i>) within the subcanopy. The understory and groundcover consisted of Virginia Creeper, Stinging Nettle (<i>Urtica dioica</i>), and Philadelphia Fleabane (<i>Erigeron philadelphicus</i>). This community has a local rank of L5 indicating this community is common and secure within the region.
CUW1-b Exotic Successional Woodland Local Rank: L+	This community was characterized by Norway Maple, Manitoba Maple and Red Pine within the sparse canopy. The understory and groundcover contained Jewelweed (<i>Impatiens capensis</i>), Red Raspberry, Gray Dogwood (<i>Cornus racemosa</i>), and Goldenrod species. This community has a local rank of L+ indicating this community is dominated by non-native species within the canopy.

Ecological Land Classification Community Type	Community Description
MEM Mixed Meadow	This community was characterized by grasses and graminoid species. There were sporadic tree species that had been planted and shrubs along the fenceline. Abundant species included Smooth Brome Grass, White Sweetclover, Wild Carrot, Creeping Thistle, and Purple Crownvetch. This community does not have a local ranking.
OAO1-T Turbid Open Aquatic Local Rank:-L+	This community included the open channel of Little Etobicoke Creek.



- Natural Heritage Study Area
- Watercourse
- Flow Direction
- Highway
- Road
- Dixie Road Bridge
- Butternut
- Ecological Land Classification**
- OAO1-T | Turbid Open Aquatic
- CUP2-A | Restoration Mixed Plantation
- CUW1-A3 | Native Deciduous Successional Woodland
- CUW1-B | Exotic Successional Woodland
- FOD7-2 | Fresh-Moist Ash Deciduous Forest
- FOD7-3 | Fresh-Moist Willow Lowland Deciduous Forest
- FOD7-a | Fresh-Moist Manitoba Maple Lowland Deciduous Forest
- FOD7-b | Fresh-Moist Norway Maple Lowland Deciduous Forest
- MEM | Mixed Meadow



City of Mississauga
Dixie-Dundas Flood Mitigation

Ecological Land Classification

Date:	November 2023	Project:	24603	Submitter:	E. Wilkinson	Reviewer:	R. Leppington
-------	---------------	----------	-------	------------	--------------	-----------	---------------

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.

I:\City\Mississauga\24603\FiguresandTables\ECL2019\Report\Figure-3\Ecological_Land_Classification.mxd - Tabbed_L - 18-Nov-23, 10:13 AM - sylvan - T10905

5.2.2 Flora

A total of 118 vascular plant species were identified within the study area through the botanical inventories and the TRCA field data from 2018. Of the 118 species identified, 58 (49.15%) species are considered native to the City of Toronto, 58 (49.15%) are considered non-native, and 2 (1.69%) was unclassified (Table 5). This high proportion of non-native species coincides with results from the Mississauga NAS that found that 43% to 52% of the species present within the study area were introduced (City of Mississauga, 2020). A full list of vegetation species found during the background review and field surveys can be found in Appendix A.

The majority of the native species (81%) were ranked as S5 (secure in Ontario). One SAR (Butternut) was observed within the study area. SAR species, as well as additional SAR flora potential, are discussed in Section 6.6.

A total of five species identified within the study area are considered locally rare (L3 ranked) and include Balsam Fir (*Abies balsamea*), Butternut (*Juglans cinerea*), Ninebark (*Physocarpus opulifolius*), Red Pine (*Pinus resinosa*), Slippery Elm (*Ulmus rubra*), and White Spruce (*Picea glauca*). An additional 15 species were identified within the study area that are ranked locally as L4 which means they are of conservation concern in the urban matrix. One Honey Locust (*Gleditsia triacanthos*) was observed. Naturally occurring Honey Locusts have an S2 designation and are classified as SCC, however, planted specimens receive no protections. Since the observed individual was a thornless variety, it is assumed to have been planted and does not qualify as SCC.

5.2.2.1 Floristic Quality Assessment

Floristic Quality Index (FQI) is a weighted species richness estimate used to compare natural areas in order to evaluate their conservation value. FQI is calculated by multiplying the mean Coefficient of Conservatism (CC) by the square root of the total number of native species present within a natural area (Oldham et al. 1995). FQI results in a number that represents the relative level of floristic quality for a given area and can be compared to FQIs of other areas.

The FQI score is 28.23, which indicates a moderate floristic quality. The quality of the native components of the study area as a whole was moderate, suggesting the study area contains mainly species that are adapted to a wide range of disturbances such as changes in hydrology, light levels, and nutrients (Table 5). The mean CC is 3.71, mean CC values that are under four indicate that the site is primarily vegetated with adaptable species that can withstand a variety of habitat changes. Table 5 identifies the quantity of plant species within each coefficient of wetness category.

Table 5 Floristic Summary and Assessment of Results

Species Diversity		
Total Species	118	-
Native Species	58	49.15%
Exotic Species ⁽¹⁾	58	49.15%
Unknown ranking	2	1.69%
Regionally Rare Species (S1 to S3)	2	-
S1 to S3 Native Species	2	1.69%
S4 Native Species	9	7.63%
S5 Native Species	47	39.83%
SNA Non-native Species	58	49.15%
SU Species	2	1.69%
Coefficient of Conservatism (CC)		
CC Mean (native) ⁽²⁾	3.71	Medium
CC 0 to 3	23	39.66%
CC 4 to 6	32	55.17%
CC 7 to 8	3	5.17%
CC 9 to 10	0	0.00%
Floristic Quality Index (FQI)		
FQI (native) ⁽³⁾	28.23	Moderate

Notes:

(1) some species are excluded from the calculation as they are currently unranked in CC.

(2) High = >4.00; Medium = 3.30 to 3.99; Low = <3.30.

(3) High = >35; Moderate = 20 to 35; Low = 1 to 19.

5.2.3 Incidental Wildlife Observations

Incidental wildlife observations are important as they may provide species identification that was not included in any specific survey. Incidental observations were conducted as part of the study and are summarized in Appendix A.

Of the species incidentally observed, one SCC species, Eastern Wood-pewee (*Contopus virens*), and one SAR, Chimney Swift (*Chaetura pelagica*), were observed within the study area. The SCC species are discussed in Section 5.2.6 and Appendix C and SAR are discussed further in Section 6.6.

A total of three wildlife species identified within the study area are considered locally rare (L2 or L3 ranked) and include Black-and-White Warbler (*Mniotilta varia*), Golden-crowned Kinglet (*Regulus satrapa*) and Yellow-bellied Sapsucker (*Sphyrapicus varius*) (Appendix A). An additional 21 species were identified within the study area that are ranked locally as L4 which means they are of conservation concern in the urban matrix.

5.3 Aquatic Resources

Little Etobicoke Creek originates north of Highway 401 in the South Slope physiographic region and flows in a general southeast direction to its confluence with Etobicoke Creek within the Iroquois Sand Plain physiographic region just upstream of The Queensway. Dominant land use within the study area consists of industrial activity in the upper subwatershed and low-medium density residential developments in the lower subwatershed. The Etobicoke Creek Aquatic Ecosystem Report (TRCA 2006) identifies Little Etobicoke Creek as a second order stream with an approximate length of 16.0 km. It has assigned a habitat classification of “small riverine warmwater” for the entire length of Little Etobicoke Creek (TRCA 2006). Small riverine warmwater systems have low mean baseflow, fluctuating water temperatures, drainage areas less than 10 km², and typically have fish communities dominated by cyprinids, Johnny Darter, and Fantail Darter (TRCA 2006).

5.3.1 Aquatic Habitat

5.3.1.1 *Upstream of Dixie Road*


Upstream of Dixie Road for approximately 500 m, Little Etobicoke Creek flowed as a defined watercourse with a slight winding pattern within a narrow natural corridor through a highly urbanized environment. Both banks contained a narrow band of mature deciduous forest approximately 8 to 20 m wide along the survey reach. Within the study area, the channel was considered entrenched with a clearly defined stream bed and an abrupt floodplain slope break. Both banks of the channel for the entire survey reach were considered stable and lined with large armour stones stacked three or four high in a slight step pattern which made it difficult to establish bankfull within the channel. The channel was densely shaded by large overhead deciduous trees and overhanging shrubs in the understory. Several large fallen trees crossed the channel resting on the armour stones along top of bank throughout the reach (Figure 5a).

A paved pedestrian path/trail was present along both banks and crossed the channel via two wooden pedestrian walking bridges approximately 200 m and 400 m upstream of Dixie Road. Approximately 30 m upstream of the first pedestrian bridge (230 m upstream of Dixie Road), a small side channel approximately 10 m long originating from a stormwater outfall contributed to the watercourse channel from the north bank. A second stormwater outfall channel joined the watercourse from the south bank immediately downstream of the second pedestrian bridge.

Channel morphology within the reach was fairly homogenous and consisted of long slow run sections interspersed with short riffles and sporadic shallow mid-channel pools. Several riffles between Dixie Road and the first pedestrian bridge were created by small bedrock drops in the stream bed. Run sections had a mean depth of 0.4 m with a wetted width of 6.0 m. Substrates in runs consisted mainly of gravel, sand, and cobble overlying bedrock. Riffles had a mean depth of 0.15 m with an average wetted width of 7.2 m. Substrates in riffles were dominated by bedrock, boulder, and cobble with pockets of gravel. Pool areas had a mean depth of 0.60 m with an average wetted width of 8.0 m. Substrates in pools were primarily gravel, sand, and cobble overlying bedrock. Bankfull measurements for all morphological features were not collected due to the presence of armour stones along the bank which made it difficult to establish true bankfull elevations within the reach.

Riparian vegetation within the study area consisted primarily of large deciduous trees and shrubs along both banks with grasses and herbaceous plants for ground cover. No instream vegetation was observed within the channel.

Habitat within the reach was limited and included cover provided by boulders and large cobbles and crevices in the bedrock and armour stone along the bank. Small amounts of overhanging woody debris and overhanging vegetation provided additional habitat opportunities. Suitable habitat to complete the various life cycle processes for the warm water bait/forage fish species present was observed throughout the reach. Gravid female Creek Chub and Blacknose Dace exhibiting spawning colouration were captured during an electrofishing survey (details in Section 5.3.2) indicating that the reach provides rearing/nursery habitat and probable spawning habitat for these species.




- Warm water Bait/Forage Fish Habitat
- Natural Heritage Study Area
- Pool
- Riffle
- Watercourse
- Flow Direction
- Highway
- Road
- Armourstone/Gabion Basket
- Fallen Tree
- Small Bedrock Knickpoint
- Woody Debris
- Pedestrian Bridge
- Run
- Field Note
- Dixie Road Bridge
- Outfall

Notes:

1. Pool with a wetted width of 5.6, wetted depth of 0.6m. Armour stone banks, with overhanging deciduous trees. Gravel, sand, and cobble substrate
2. Riffle with a wetted width of 7.2m and wetted depth of 0.15m. Armourstone banks with overhanging deciduous trees. Cobble, boulders and gravel substrate.
3. Run with a wetted width of 8m and wetted depth of 0.30m. Armourstone banks with overhanging deciduous trees. Gravel, sand and cobble substrate.

General Comments:

- Both banks lined with large armour stone stacked four stones high in slight step pattern.
- Informal trails through deciduous forest areas on both banks.
- Thin layer of substrates overlying bedrock throughout survey reach
- Channel well shaded by overhanging shrubs and trees on both banks.



Reference: Contains information licensed under the Open Government Licence - Ontario. Contains data obtained by Matrix Solutions Inc. and the City of Mississauga. Imagery: Service Layer Credits: © 2023 Microsoft Corporation © 2023 Maxar © CNES (2023) Distribution Airbus DS

1:2,000 metres
20 0 20 40
NAD 1983 UTM Zone 17N



City of Mississauga
Dixie-Dundas Flood Mitigation

**Fish Habitat
Upstream of Dixie Road**

Date: November 2023 Project: 24603 Submitter: E. Wilkinson Reviewer: R. Leppington

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.

I:\CityOfMississauga\24603\FiguresandTables\ECL\2019\Report\Figure-5a-Fish_Habitat_Upstream_of_Dixie_Road.mxd - Tabbed_L - 16-Nov-23, 10:32 AM - sjlam - TID005

5.3.1.2 *Downstream of Dixie Road*

Similar to the section upstream of Dixie Road, the watercourse continued as a defined channel within a narrow natural corridor downstream of Dixie Road for approximately 700 m. The forested area along the banks was wider and varied from 25 to 55 m wide. The channel was densely shaded by large overhead deciduous trees and overhanging shrubs in the understory. Both banks of the channel were lined with large armour stones in a slight step pattern to provide bank stabilization during high flows. The armour stone wall continues for approximately 500 m downstream of Dixie Road, leaving the remaining 200 m until the crest of the large meander bend with natural banks (Figure 5b)

A paved pedestrian trail/path was present on the north bank with several small informal trails noted with the deciduous forest. Downstream of Dixie Road, the watercourse flowed along the south bank of the channel with a large, exposed cobble bar along the north side of the channel. Approximately 550 m downstream of Dixie Road, a small side channel enters from the north which creates a large pool (1 m depth) at the confluence.

Channel morphology within the reach downstream of Dixie Road consisted of a riffle/pool sequence with long riffle sections and a few shallow-deep pools. Mean depth within riffles was 0.2 m with an average wetted width of 6.0 m. Mean depth within pools was 0.5 m with an average wetted width of 5.5 m. Substrates within the reach consisted primarily of boulder, cobble, and bedrock with pockets of gravel and sand.

Riparian vegetation consisted of overhanging deciduous trees and shrubs with herbaceous plants and grasses for ground cover. Instream aquatic macrophyte growth primarily consisted of submergent algae which covered larger substrates.

Habitat immediately downstream of Dixie Road was limited; however, further downstream there was more naturalized bank habitat which provided undercut banks, exposed root wads, and woody debris. The majority of habitat provided consisted of instream boulders and cobble and crevices in bedrock and armour stone. Existing habitat provides sufficient opportunity for the warmwater bait/forage fish community present to complete their various lifecycle requirements. Gravid female Creek Chub and Blacknose Dace exhibiting spawning colouration were captured during an electrofishing survey (details in Section 5.3.2) indicating that the reach provides rearing/nursery habitat and probable spawning habitat for these species.


Easting (m)

Northing (m)



- Warm water Bait/Forage Fish Habitat
- Natural Heritage Study Area
- Pool
- Riffle
- Watercourse
- Flow Direction
- Highway
- Road
- Armourstone/Gabion Basket
- Fallen Tree
- Woody Debris
- Run
- Field Note
- Dixie Road Bridge
- Outfall
- Sand Bar
- Cobble Bar
- Undercut

Notes:
 4. Pool with a wetted width of 6m and a wetted depth of 0.52m. Armourstone banks with overhanging deciduous trees. Substrate comprised of cobble, gravel with sand overlay and areas of exposed bedrock.
 5. Riffle with a wetted width of 6m and wetted depth of 0.18m. Armourstone banks with overhanging deciduous trees. Substrate comprised of cobble, boulder and exposed bedrock.
 6. Pool with a wetted width of 4m and depth of 0.50m. Armourstone ends and banks are natural with no armouring. Substrate is sand and exposed clay.



Reference: Contains information licensed under the Open Government Licence - Ontario. Contains data obtained by Matrix Solutions Inc. and the City of Mississauga. Imagery: Service Layer Credits: © 2023 Microsoft Corporation © 2023 Maxar © CNES (2023) Distribution Airbus DS



City of Mississauga
Dixie-Dundas Flood Mitigation

Fish Habitat Downstream of Dixie Road

Date: November 2023 Project: 24603 Submitter: E. Wilkinson Reviewer: R. Leppington

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.

5.3.1.3 *Upstream of Dundas Street East*

The section of channel upstream of Dundas Street for approximately 500 m had a mix of bank treatments which included large boulders at the crest of the meander bend, gabion baskets at the downstream extent of the bend, armour stone at the most downstream extent, and intermittent naturalized banks. Overall, the area was shaded by overhanging deciduous trees; however, the forested riparian community narrowed from an approximate width of 30 m close to Dixie Road to a 5 m width approaching Dundas Street (Figure 5c).

The crest of the large meander bend contained a riffle feature with large circular cobbles and boulders within the channel. A large sand bar developed downstream of the riffle into a small pooled habitat. The pools contained large substrates such as cobbles and boulders, with an overlay of silt and sand. Substrate within the riffles consisted primarily of cobbles, boulders, and gravel with areas of exposed bedrock. The armouring of the banks changed to gabion baskets which created a long deep pooled system with depths of 0.70 to 0.80 m deep and 5 m wetted width for approximately 50 m. Naturalized banks were present moving downstream toward Dundas Street. The naturalized banks had areas of undercutting (0.26 m), wood debris, and a mixture of pool, riffle and run habitats. The riffle and run habitats ranged between 0.15 to 0.26 m wetted depths and consisted primarily of cobbles, boulders, and gravel with areas of exposed bedrock. The channel began to narrow with a wetted width of 3 m and armour stone along the west bank. The presence of armour stone increased approaching Dundas Street with a large pool (0.88 m deep).

The majority of habitat provided consisted of instream boulders and cobble, and crevices in bedrock and armour stone. In some of the more naturalized area, there were habitat features such as undercut banks, root wads, and wood debris. The results of the electrofishing survey (details in Section 5.3.2) indicated the presence of a warmwater baitfish community. Existing habitat provides opportunity for the warmwater bait/forage fish community present to complete their various lifecycle requirements. Throughout the reach there were schools of young of the year fish which were observed within the sandy pooled habitats. This indicates that this reach provides rearing/nursery habitat and confirmed spawning habitat for these species.



- Warm water Bait/Forage Fish Habitat
- Natural Heritage Study Area
- Pool
- Riffle
- Watercourse
- Highway
- Road
- Armourstone/Gabion Basket
- Fallen Tree
- Woody Debris
- Run
- Field Note
- Outfall
- Sandbar
- Cobble Bar
- Undercut
- Overhanging Banks
- Vertical Bank

Notes:

7. Riffle with wetted width of 2m and wetted depth of 0.16m. Right bank has vertical banks, Left bank has large cobble bar. Confluence of tributary. Woody debris in centre channel at the end of riffle.

8. Pool with wetted width of 5m and wetted depth of 0.82m. Boulders/cobble with sand and silt overlay. Falling gabion baskets along the banks.

9. Run with etted eidth of 5m and wetted depth of 0.43m. Left bank has under cutting with woody debris along right bank. Substrate contains large cobble, boulders and exposed bedrock.

10. Pool with 5m wetted width and 0.88m wetted depth. Left bank is vertical and bare. Substrate is silt, sand and cobble.



Reference: Contains information licensed under the Open Government Licence - Ontario. Contains data obtained by Matrix Solutions Inc. and the City of Mississauga. Imagery: Service Layer Credits: © 2023 Microsoft Corporation © 2023 Maxar © CNES (2023) Distribution Airbus DS



City of Mississauga
Dixie-Dundas Flood Mitigation

Fish Habitat Upstream of Dundas Street

Date: November 2023 Project: 24603 Submitter: E. Wilkinson Reviewer: R. Leppington

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.

I:\City\Mississauga\24603\FiguresandTables\ECL\2019\Report\Figure-5c-Fish_Habitat_Upstream_of_Dundas_Street.mxd - Tabloid_L - 16-Nov-23, 10:41 AM - slyam - TID005

5.3.1.4 Downstream of Dundas Street East


The section of the channel downstream of Dundas Street East for approximately 360 m had a mix of bank treatments which included armour stone and gabion baskets at the most upstream extent near Dundas Street East, intermittent naturalized banks with terrace formations, or steeply eroded banks. Overall, the area was partially shaded by overhanging deciduous trees (Figure 5d).

Channel morphology of the stream consisted of a series of riffles, pools, and runs. Riffles averaged a depth of 5 to 40 cm deep, with a wetted width of 1 to 4 m wide, gradually narrowing further downstream. Pools averaged a depth of 0.8 to 1.5 m with a wetted width of 5 to 9.5 m. Runs averaged a depth of 15 to 50 cm with a wetted width of 6 to 9.5 m. Substrates varied within the reach consisting of large cobble, cobble, and some sand and gravel or mostly cobble with sand and gravel.

Starting at the Dundas Street East culvert there was a run that featured large flat rocks covered in filamentous algae with cobbles in between. Further downstream at the end of the armour stone headwalls the channel transitioned into a riffle feature with large cobble consisting of natural stone and concrete blocks, cobble, and some sand and gravel. After a 30 to 40 cm drop, there was a pool that was completely shaded under an overhanging tree that had fallen across the waterway. The right bank was lined with gabions, the ones submerged in the water had been washed out creating an artificial undercut bank (0.30 m) that transitioned from a run into a pool that was concentrated on the right bank along the gabions and back into a run. At the end of the gabions the left bank became very steep and undercut (0.28 m) with woody debris, while the right bank was a cobble bar with overhanging vegetation. This was followed by a pool that was 80 cm deep that switched to a run containing some metal debris. The next riffle had a gentler slope followed by a pool with terraced slopes on the right bank. The left bank contained woody and metal debris and exposed roots. Moving further downstream through another riffle and pool there was more debris on the right bank with evidence of filter cloth on the left bank. The last riffle was near the SWM outlet on the left bank with the right bank being heavily eroded, exposing soil and tree roots. The last pool had undercut banks and slumping or evidence of slumping with exposed tree roots.

Riparian vegetation consisted of overhanging deciduous trees and shrubs with herbaceous plants and grasses for ground over. Filamentous algae was present on some rocks within the stream.

The majority of habitat provided consisted of instream boulders and cobble, washed out and undercut gabion cages, overhanging fallen trees as well as multiple sections containing woody debris with some undercut banks and exposed roots. The results of the electrofishing survey (details in Section 5.3.2) indicated the presence of a warmwater baitfish community. Existing habitat provides opportunity for the warmwater bait/forage fish community present to complete their various lifecycle requirements. There was a Blacknose Dace with spawning colouration captured during the electrofishing survey indicating that the reach provides rearing/nursery habitat and probable spawning habitat for this species.




- Natural Heritage Study Area
- Warm water Bait/Forage Fish Habitat
- Outfall
- Pool
- Riffle
- Watercourse
- Highway
- Road
- Run
- Armourstone/Gabion Basket
- Fallen Tree
- Small Cinderblock Retaining Wall
- Woody Debris
- Field Note
- Large Rock
- Cobble Bar
- Undercut
- Vertical Bank

Notes:

1. Pool with wetted width of 8m and wetted depth of 1m approximately. Right bank is lined with failing gabions, left bank slopes gradually with overhanging deciduous vegetation. Mostly sandy gravel with 10% cobble substrate underlined with clay.
2. Run with a wetted width of 9.5m and wetted depth of 15cm approximately. Right bank slopes gently, left bank is steeper with exposed roots. Both banks are heavily vegetated. Cobble, sand, gravel substrate.
3. Riffle with wetted width of 4m and wetted depth of 10cm approximately. Right bank heavily eroded with exposed soil and roots, left bank eroded due to an outlet. Cobble substrate with minimal large cobble and small gravel.

General Comments:

- Armour stone present closer to the Dundas St E overpass with failing gabions on the left bank for a portion of the reach.
- Right bank generally slopes more gently with the left bank being steeper and more eroded.
- Informal trails through deciduous forest areas on both banks.
- Channel generally shaded by overhanging shrubs and trees on both banks.



Reference: Contains information licensed under the Open Government Licence - Ontario. Contains data obtained by Matrix Solutions Inc. and the City of Mississauga. Imagery: Service Layer Credits: © 2023 Microsoft Corporation © 2023 Maxar © CNES (2023) Distribution Airbus DS

1:1,200 metres
10 0 10 20
NAD 1983 UTM Zone 17N



City of Mississauga
Dixie-Dundas Flood Mitigation

Fish Habitat Downstream of Dixie Road

Date: November 2023 Project: 24603 Submitter: E. Wilkinson Reviewer: R. Leppington

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.

I:\City\Mississauga\24603\FiguresandTables\ECL\2019\Report\Figure-5d\Fish_Habitat_Downstream_of_Dundas_Street.mxd - Tabloid_L - 16-Nov-23, 10:12 AM - s.yam - TID005

5.3.2 Fish Community

The background review identified historical fisheries information for Little Etobicoke Creek from 1946 to 1991 and more recent records in the Lower Etobicoke Creek from 2001 to 2016 (TRCA 2006; TRCA 2019b) approximately 1.1 km downstream of Dundas Street (Table 6). The historical fisheries data indicates that presence of various common warmwater and cool water species, with the exception of the Redside Dace (*Clinostomus elongatus*), which is listed as Endangered both provincially and federally. However, the record of Redside Dace is considered historical, as it has not been observed in Little Etobicoke Creek since 1946.

Table 6 Historical Fisheries Data at Little Etobicoke Creek Within and Surrounding the Study Area

Common Name	Scientific Name	SARA ⁽¹⁾	ESA ⁽²⁾	Little Etobicoke 1946-1991	Lower Etobicoke 2001-2016
Eastern Blacknose Dace	<i>Rhinichthys atratulus</i>	-	-	X	X
Bluntnose Minnow	<i>Pimephales notatus</i>	-	-	X	X
Brook Stickleback	<i>Culaea inconstans</i>	-	-	X	-
Central Stoneroller	<i>Campostoma anomalum</i>				X
Common Shiner	<i>Luxilus cornutus</i>	-	-	X	X
Creek Chub	<i>Semotilus atromaculatus</i>	-	-	X	X
Fathead Minnow	<i>Pimephales promelas</i>	-	-	X	-
Golden Shiner	<i>Notemigonus crysoleucas</i>	-	-	-	X
Johnny Darter	<i>Etheostoma nigrum</i>	-	-	X	X
Longnose Dace	<i>Rhinichthys cataractae</i>	-	-	X	X
Rainbow Darter	<i>Etheostoma caeruleum</i>	-	-	X	-
Redside Dace	<i>Clinostomus elongatus</i>	Endangered (Schedule 1)	Endangered	X	-
Rock Bass	<i>Ambloplites rupestris</i>	-	-	X	-
White Sucker	<i>Catostomus commersonii</i>	-	-	X	X

Notes:

(1) Species at Risk Act (SARA; Government of Canada 2021)

(2) Endangered Species Act, 2007 (ESA; Government of Ontario 2021a)

In 2019, Matrix conducted an electrofishing survey at four stations throughout the Little Etobicoke Creek study area (Figure 2). The stations were located 200 m upstream of Dixie Road (LEC1), 50 m and 400 m downstream of Dixie Road (LEC2 and LEC3), and 200 m upstream of Dundas Street (LEC4). In 2021, Matrix conducted an electrofishing survey at one additional station downstream of Dundas Street East (LEC5).

The station was located approximately 250 m downstream of the culvert, with a total of 60 m of creek surveyed. A total of 85 fish across 7 species were caught in across the 2019 and 2021 fish community surveys (Table 7). All the species captured are considered to be common, tolerant, warmwater/coolwater species within Ontario.

Table 7 Fish Community Results for Little Etobicoke Creek in 2019 and 2021

Species Common Name	LEC1	LEC2	LEC3	LEC4	LEC5 ⁽¹⁾
Central Stoneroller	-	-	1	-	-
Creek Chub	8	8	1	-	-
Eastern Blacknose Dace	3	1	5	10	2
Longnose Dace	1	3	19	16	4
White Sucker	1	1	-	-	-
Brown Trout	-	-	-	-	1
Total	13	13	26	26	7

Notes:

(1) Survey conducted in 2021

6 SIGNIFICANT NATURAL HERITAGE FEATURES AND FUNCTIONS

Significant natural heritage features and functions include those listed in the Provincial Policy Statement (MMAH 2020), and the NHRM (MNR 2010). Reference was also obtained from the natural heritage system from the City’s Official Plan (City of Mississauga 2021). The findings of the site investigations were cross-referenced with the criteria provided in these documents in order to identify the confirmed or potential presence of significant natural heritage features.

The following significant features were not present within 120 m of the study area:

- ANSIs
- Environmentally Significant Areas
- Provincially Significant Wetlands
- Wetlands or Unevaluated Wetlands

Significant features that are present within the study area are discussed further in Sections 6.1 to 6.6.

6.1 Significant Valleylands and Corridors

Valleylands are linear natural areas that occur in a valley or other landform depressions that have water flowing through or standing for some period of the year (MNR 2010). These areas are important corridors which provide unique features and functions to an area as well as linkages between terrestrial and aquatic habitats.

Little Etobicoke Creek and associated valley lands are a part of the City's "Significant Natural Areas" (City of Mississauga 2021). The overall function of the adjacent natural lands associated with Little Etobicoke Creek play a meaningful role in the urban landscape and it is considered a significant valleyland.

6.2 Significant Woodlands

Section 6.3.12 of the City's Official Plan (City of Mississauga 2021) states the criteria needed to meet the significant woodlands designation within the City of Mississauga. Since the woodland surrounding Little Etobicoke Creek is a continuous system which is greater than 0.5 ha and is within 30 m of a watercourse, it is considered a significant woodland (City of Mississauga 2021).

6.3 Fish and Fish Habitat

As presented in Section 5.3, the study area does contain fish as well as permanent fish habitat within Little Etobicoke Creek.

Fish and fish habitat are regulated by Fisheries and Oceans Canada (DFO) under the Fisheries Act (Government of Canada 2019). The Fisheries Act requires that projects avoid causing the death of a fish or a harmful alteration, disruption or destruction (HADD) of fish habitat unless authorized by the Minister or a designated representative (Government of Canada 2019). The determination of death of fish or HADD is typically done through a self-assessment process.

6.4 Linkages and Corridors

Linkages and corridors are important features within a natural system. These features are continuous, often linear bands of vegetation in the landscape which provide opportunities to connect natural areas and provide cover for wildlife movement and dispersal of otherwise isolated populations.

Little Etobicoke Creek has been designated as a significant valleyland. This area represents a significant linkage for both terrestrial and aquatic organisms. The wooded riparian area along the edge of the creek provides a linkage to other natural areas within the system.

6.5 Species at Risk

A total of 17 SAR and 14 SCC were identified as having records occurring within or in the vicinity of the study area during the Matrix background review. This number was refined based on habitat characteristics present within the study area as observed through field investigations. The screening determined there is a moderate probability for six SAR and four SCC, and a confirmed presence of two SAR and one SCC. The list below provides a description of the species with confirmed or moderate probability of occurrence within the study area. The full screening can be found in Appendix C.

6.5.1 Confirmed SAR

Butternut: A Butternut was observed within the study area. This species is designated as Endangered under the ESA and SARA, however only the ESA would apply in the case of this project. The protection under the ESA only applies to pure Butternut individuals. To identify whether a Butternut is a pure species or a hybrid it must be assessed by a Butternut Health Assessor. If the tree is identified as a pure species and works will occur within 25 m of the tree, then the project will need to be registered under the MECP Notice of Activity.

Chimney Swift: this species is designated as Threatened under ESA and SARA. This species utilizes chimneys for nesting and roosting during the breeding season. The ESA general habitat protection identifies chimneys used for roosting and nesting purposes (Government of Ontario 2021a). Within the study area, Chimney Swifts were observed flying over the study area foraging for insects. Although this species was recorded within the study area, there were no suitable nesting chimneys observed.

6.5.2 Moderate Probability SAR

SAR Bats [Tri-colored Bat (*Perimyotis subflavus*); Northern Myotis (*Myotis septentrionalis*); Little Brown Myotis (*Myotis lucifugus*)]: The forested ravine habitat (primarily east of Dixie Road and east of Dundas Street East) has been identified as potential suitable habitat for SAR bat species. These species and their habitats are protected under the ESA. In order to confirm the presence or absence of these SAR bat species utilizing the potential habitat identified in Section 5.2.4.1 and depicted in Figure 5, an acoustic survey is required following the Survey Protocol for Species at Risk Bats within Treed Habitats: Little Brown Myotis, Northern Myotis & Tri-Colored Bat (MNRF 2017). Acoustic surveys would only be required if a tree or forested area identified as potential suitable habitat would be impacted or removed as part of project construction. If the potential SAR habitat is occupied by SAR bats, the project will need approval or permitting from the MECP for the removal of any confirmed SAR bat habitat.

Bank Swallow (*Riparia riparia*): Bank swallow is listed as Threatened both provincially and federally. This species can be found breeding in vertical faces such as lake and riverbanks, soil stockpiles, and sand and gravel pits. Foraging habitat consists of open areas such as meadows, pastures, and grasslands. Candidate habitat exists along any eroded vertical banks within Little Etobicoke Creek.

Eastern Whip-poor-will (*Antrostomus vociferus*): Eastern Whip-poor-will is listed as Threatened both provincially and federally. This species can be found in areas that contain a mix of open and forested habitat such as open woodlands, forests with openings, and savannahs. Candidate habitat exists within the FOD communities within the study area that are surrounded by open meadow.

Red-headed Woodpecker (*Melanerpes erythrocephalus*) – Red-headed Woodpecker is listed as Endangered provincially and threatened federally. This species can be found in woodlands and woodland edges, including oak and beech forests, grasslands, orchards, riparian forests, beaver ponds, burns, parks,

golf courses, and cemeteries. Dead trees are used for nesting and perching. Candidate habitat exists within the FOD communities within the study area.

6.5.3 Confirmed SCC

Eastern Wood-pewee (*Contopus virens*) – Eastern Wood-pewee is listed as Special Concern both provincially and federally. This species can be found in the mid-canopy layer of forest clearings, edges of deciduous and mixed forests, early successional clearings, coniferous forests, mixed forests, deciduous forests, deciduous swamps, mixed swamps, and cultural woodlands. An eastern wood-pewee was heard vocalizing during the 2019 field visits, suitable habitat was present within the FOD communities within the study area.

6.5.4 Moderate Probability SCC

Canada Warbler (*Cardellina canadensis*) – Canada Warbler is listed as Special Concern both provincially and federally. This species can be found in deciduous or coniferous forests with a well-developed, dense shrub layer; commonly in wet or riparian areas. This may also include stands that are regenerating after natural disturbances. Candidate habitat exists within the FOD communities within the study area.

Wood Thrush (*Hylocichla mustelina*) – Wood Thrush is listed as Special Concern provincially and Threatened federally. This species can be found in mature deciduous and mixed forests, or in moist stands of trees with well-developed undergrowth. Their nests are built in live saplings, trees, or shrubs, especially sugar maple or American beech. Candidate habitat exists within the FOD communities within the study area.

Barn Swallow (*Hirundo rustica*) – Barn Swallow is listed as Special Concern both provincially and federally. This species constructs its cup-shaped mud nests on rough-cut wood surfaces on human-made structures such as open barns, under bridges, and in culverts. This species forages in grassy fields, pastures, cropland, lake and river shorelines, cottage areas, farmyards, islands, wetlands, and tundra. Candidate nesting habitat exists along the bridges and culverts within the study area, candidate foraging habitat exists along the creek.

Monarch (*Danaus plexippus*) – Monarch is listed as Special Concern both provincially and federally. This species is found in open or disturbed habitats such as roadsides, fields, wetlands, prairies, and open forests where milkweed is present. Milkweed was observed within the study area during field visits, candidate habitat exists wherever milkweed is present.

6.5.5 Bat Maternity Roosting Survey

The Tri-colored Bat and the two Myotis species require different roosting habitat characteristics. Little Brown Myotis and Northern Myotis roost in-tree cavities, crevices, and exfoliating bark in wooded areas located near water. The Tri-colored Bat most often roosts in foliage (both dead and alive) within or

below the canopy. Often, Oak (*Quercus*) species are utilized for roosting because the leaves are retained longer in the fall season; however, Maple (*Acer*) species are also used. Tri-colored Bats forage along riparian corridors and open water.

Identifying suitable roost trees for Little Brown Myotis and Northern Myotis included recording the locations and decay classes of all snags that exhibited appropriate attributes including cavities, loose bark, cracks, or knot holes. Decay class is an important consideration when ranking the quality of roosting habitat, with trees in early stages of decay typically being of higher quality than those in later stages of decay. Each snag tree was ranked into a decay class of 1 through 6. Class 1 has the least amount of decay present, and class 6 has the most. Class 1-3 trees are typically preferred for roosting because their bark and wood are still mostly intact, offering greater protection from the elements compared to a more decayed tree that has lost all its bark and whose wood has been decomposing for many years. Table 8 summarizes the suitable maternity roost trees for Little Brown Myotis and Northern Myotis throughout the entire study area.

Table 8 Summary of Little Brown Myotis and Northern Myotis Suitable Roost Trees within Study Area (Leaf-off Survey)

Tree Number	Tree Species	Diameter at Breast Height (DBH)	Decay Class
1	unknown species	28	5
2	Maple sp.	35	4
3	Willow sp.	50	2
4	Willow sp.	55	5
5	Willow sp.	50	5
6	Willow sp.	55	2
7	Willow sp.	60	4
8	Oak sp.	24	2
9	unknown species	22	4
10	Maple sp.	40	2
11	Maple sp.	37	2
12	Maple sp.	20/20	5
13	Maple sp.	25	4
14	Maple sp.	30	4
15	unknown species	20	4
16	Maple sp.	50/40	2
17	unknown species	24	4
18	Ash sp.	15/15/15	5
19	Willow sp.	65	1
21	Willow sp.	50/45	2
22	Willow sp.	70/70/60	2
25	Eastern Cottonwood	70/70	4

Tree Number	Tree Species	Diameter at Breast Height (DBH)	Decay Class
26	unknown sp.	30	6
27	Ash sp.	30	3
28	Eastern Cottonwood	50	3
29	unknown sp.	32	3
30	Ash sp.	15	3
31	Maple sp.	20	4
32	Eastern Cottonwood	50/25	5

Notes:

Trees in bold indicate high-quality snag trees.

A total of 34 snags greater than 10 cm DBH were recorded during the snag evaluation, of which 29 are within the Dixie-Dundas study area. 13 of the 29 trees located within the study area are considered high-quality snags.

Identifying suitable roost habitat for Tri-colored Bats included recording the location of Oak trees greater than 10 cm DBH, Maple trees greater than 10 cm DBH if they included dead leaf clusters, and Maple trees greater than 25 cm DBH (dead leaf cluster or not). Table 9 summarizes the suitable maternity roost trees for Tri-Colored bats throughout the entire study area.


Table 9 Summary of Tri-Colored Bat Suitable Roost Trees within Study Area (Leaf-on Survey)

Tree Number	Tree Species	Diameter at Breast Height (DBH)	Tree Status	Dead/Dying Leaf Cluster
1	Red Oak	44	Live	No
2	Red Oak	36	Live	No
3	Red Oak	41	Live	No
4	Red Oak	54/50	Live	No
5	Red Oak	51	Live	No
6	White Oak	13	Live	No

A total of six large diameter oak trees were recorded within the study area. In addition to the oak species specified in Table 9, there was a large quantity of maple trees (Manitoba maple [*Acer negundo*] and Norway maple [*Acer platanoides*]) throughout the study area that were greater than 25 cm DBH. No maple trees were recorded with visible dead leaf clusters. Although maple species are usually only recorded if Oak species are absent from the study area the oaks recorded were found in a central location downstream of Dixie Road; therefore, maple species were assessed as well. Maple trees were located in six of the eight vegetation community types. The 25 high-quality snag trees, oak trees, and ELC polygons containing maples greater than 25 cm DBH are considered potential SAR bat habitat and have been mapped to avoid their removal where possible (Figure 5).



- Natural Heritage Study Area
 - Area with Maples >25 cm DBH
 - Watercourse
 - Flow Direction
 - Highway
 - Road
 - Dixie Road Bridge
 - Butternut
 - Leaf-On Assessment - Oaks and Maples
 - Leaf-Off Assessment - Snags
- Ecological Land Classification**
- OAO1-T | Turbid Open Aquatic
 - CUP2-A | Restoration Mixed Plantation
 - CUW1-A3 | Native Deciduous Successional Woodland
 - CUW1-B | Exotic Successional Woodland
 - FOD7-2 | Fresh-Moist Ash Deciduous Forest
 - FOD7-3 | Fresh-Moist Willow Lowland Deciduous Forest
 - FOD7-a | Fresh-Moist Manitoba Maple Lowland Deciduous Forest
 - FOD7-b | Fresh-Moist Norway Maple Lowland Deciduous Forest
 - MEM | Mixed Meadow



Reference: Contains information licensed under the Open Government Licence - Ontario. Contains data obtained by Matrix Solutions Inc. and the City of Mississauga. Imagery (2018) provided by the City of Mississauga used under license.



City of Mississauga
Dixie-Dundas Flood Mitigation

Potential Bat Maternity Roosting Habitat


Date: November 2023 Project: 24603 Submitter: E. Wilkinson Reviewer: R. Leppington

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.


6.6 Significant Features and Functions Summary

Based on the background review and site investigations to date, the potential and confirmed significant features and functions that are present within the study area are summarized below and depicted in Figure 6:

- Significant woodland (confirmed)
- Significant valleyland (confirmed)
- Fish and fish habitat (confirmed)
- Confirmed SAR and SCC:
 - ✦ Butternut (SAR)
 - ✦ Chimney Swift (SAR)
 - ✦ Eastern Wood-pewee (SCC)
- Moderate probability of occurrence for six SAR and four SCC



- Natural Heritage Study Area
- TRCA Regulation Limit
- Significant Natural Areas and Natural Green Spaces
- Watercourse
- Flow Direction
- Highway
- Road
- Potential SAR Bat Habitat - Snag
- Potential SAR Bat Habitat - Oak Tree
- Potential SAR Bat Habitat - Maple Trees



Reference: Contains information licensed under the Open Government Licence - Ontario. Contains data obtained by Matrix Solutions Inc. and the City of Mississauga. Imagery: Service Layer Credits: © 2024 Microsoft Corporation © 2024 Maxar © CNES (2024) Distribution Airbus DS



City of Mississauga
Dixie-Dundas Flood Mitigation

Natural Heritage Features and Buffers

Date:	March 2024	Project:	24603	Submitter:	E. Wilkinson	Reviewer:	R. Leppington
-------	------------	----------	-------	------------	--------------	-----------	---------------

Disclaimer: The information contained herein may be compiled from numerous third party materials that are subject to periodic change without prior notification. While every effort has been made by Matrix Solutions Inc. to ensure the accuracy of the information presented at the time of publication, Matrix Solutions Inc. assumes no liability for any errors, omissions, or inaccuracies in the third party material.

7 DESCRIPTION OF ALTERNATIVES

7.1 Dixie Area Alternative Solutions

Three alternative solutions were developed for the Dixie Area that represented a different approach to keep flow within the valley corridor. All alternatives would require some tree removals, although some more than others. All alternatives would also require some short-term impacts to the aquatic ecosystem during construction, but no long-term effects would occur once the watercourse has stabilized.

7.1.1 Dixie AS1: Improved Conveyance with a Minimized Footprint

Dixie AS1 centres around improving water conveyance through an oversized and incised channel. 600 m of the channel would be lowered by approximately 1 m. The channel would be widened but would not be well-connected to the floodplain. This alternative would require a new bridge span at Dixie Road of 26 m, and Dixie Road would need to be raised by approximately 1.7 m.

From an ecological perspective, Dixie AS1 would have the lowest impacts to the terrestrial environment, with 2.27 ha of treed areas being impacted to reconstruct the Dixie Road bridge and lower the channel. The existing vegetation along the valley would remain. No opportunities for recreational amenities or fish habitat improvements are present with this alternative.

7.1.2 Dixie AS2: Improved Conveyance by Making Room for Little Etobicoke Creek

Dixie AS2 centres around improving conveyance through a lowered and widened channel and a connected and lowered floodplain. 600 m of the channel would be lowered by approximately 0.5 m to create a low-flow channel. Floodplain shelves would be added to convey water when low-flow conditions are exceeded. This alternative would require two new bridge spans at Dixie Road of 45 m total length, and Dixie Road would need to be raised by approximately 0.4 m.

From an ecological perspective, Dixie AS2 would require the most tree removals along the project study area to widen the valley and channel corridor, with 3.77 ha of treed areas being impacted. The butternut tree identified within this area is not anticipated to be directly impacted; however, the 25 m surrounding buffer would be infringed upon. A mitigation plan for this tree (should the butternut health assessment show it to be a pure butternut), would be required. Should acoustic surveys identify bat presence within the study area, mitigation for the removal of SAR bat roosting trees would also be required. Given the high percentage of non-native tree species within the wooded area, the removal of vegetation along the corridor would present an opportunity to revegetate the corridor with more valuable native species. Additionally, this alternative presents the highest potential for fish habitat improvements.

7.1.3 Dixie AS3: Flood Containment with Mitigation for Upstream Impacts

Dixie AS3 centres around containing flood flows within the existing valley corridor by using a flood protection landform (FPL; a permanent massive earthen structure with an engineered clay core). The FPL

would extend 1.25 km along Little Etobicoke Creek. Minor channel widening would also occur, no channel lowering would be required. This alternative would require a new bridge span at Dixie Road of 28 m, and Dixie Road would need to be raised by approximately 2.5 m. The FPL's large footprint would also cause significant property impacts.

From an ecological perspective, Dixie AS3 would require some vegetation removals along the south bank of Little Etobicoke Creek to accommodate the FPL, with 2.38 ha of treed areas being impacted. The FPL would also likely encroach onto existing trails, requiring their relocation or complete removal. No opportunities for additional recreational opportunities or fish habitat improvements are present with this alternative.

7.1.4 Dixie Do Nothing

The do nothing alternative is a required solution as part of the Municipal Class EA process, to provide a reference comparison for the other alternative solutions. The Dixie Do Nothing approach would not cause any disturbance to the terrestrial ecosystem because there would be no construction. For the aquatic ecosystem, there would be no opportunities for fish habitat improvement, and continued erosive flows would continue to degrade the habitat quality.

7.1.5 Preferred Alternative

Overall, Dixie AS2 is the preferred alternative. Although it is the alternative with the greatest amount of tree removals, the removals provide an opportunity to revegetate the stream corridor with a selection of native species that provide greater ecological value than the existing tree community that is comprised of a large percentage of non-native species. Although this restoration will provide a long-term benefit to the community, it should be acknowledged that it will take several decades for the planted trees to grow to maturity and start providing the desired ecosystem services. In the interim, there will be a medium-term net loss as wildlife will not be able to utilize the young forest to the same degree as they presently use the mature, non-native forest. From an aquatic perspective, Dixie AS2 provides the greatest opportunity for fish habitat improvement through increased connectivity between the creek and its floodplain.

7.2 Dundas Area Alternative Solutions

Three alternative solutions were developed for the Dundas Area that represented a different approach to keep flow within the valley corridor. All alternatives would require tree removals, although AS1 and AS3 would require more than AS2.

7.2.1 Dundas AS1

Dundas AS1 includes a 25 m single-span bridge at Dundas Street with downstream floodplain conveyance improvements. Dundas Street East would need to be raised by approximately 0.75 m, resulting in a road disturbance length of 190 m. Conveyance improvements would include floodplain improvements and channel widening.

From an ecological perspective, Dundas AS1 would provide some opportunities for improved terrestrial connectivity through the downstream floodplain improvements. Tree removals would be required for the bridge replacement and channel widening (0.6 ha of impacted treed area) and for the floodplain improvements (0.94 ha of impacted treed area), impacting a total of 1.55 ha of treed area. An additional 0.84 ha of mixed meadow community would be impacted. Most potential fish habitat improvements with no piers present to provide constraints.

7.2.2 Dundas AS2

Dundas AS2 includes a 38 m bridge span at Dundas Street with no downstream floodplain conveyance improvements. Dundas Street East would need to be raised by approximately 0.5 m, resulting in a road disturbance length of 140 m.

From an ecological perspective, Dundas AS2 would provide some opportunities for improved terrestrial connectivity through the larger bridge span. AS2 would require the smallest amount of tree removals since no floodplain improvements would occur. A total of 0.6 ha of treed area would be impacted with the bridge replacement and channel widening. An additional 0.01 ha of mixed meadow community would be impacted. Some potential fish habitat improvements, piers provide a habitat constraint.

7.2.3 Dundas AS3

Dundas AS3 includes a 38 m bridge span at Dundas Street with downstream floodplain conveyance improvements. Dundas Street East would need to be raised by approximately 0.2 m, resulting in a road disturbance length of 70 m. Conveyance improvements would include floodplain improvements and channel widening.

From an ecological perspective, Dundas AS3 would provide the most opportunities for improved terrestrial connectivity through the larger bridge span and downstream floodplain improvements. AS3 would require the same amount of tree removals as AS1, with the bridge replacement and channel widening (0.6 ha of impacted treed area) and the floodplain improvements (0.94 ha of impacted treed area) impacting a total of 1.55 ha of treed area. An additional 0.84 ha of mixed meadow community would be impacted. Some potential fish habitat improvements, piers provide a habitat constraint.

7.2.4 Dundas Do Nothing

The do nothing alternative is a required solution as part of the Municipal Class EA process, to provide a reference comparison for the other alternative solutions. The Dundas Do Nothing approach would not cause any disturbance to the terrestrial ecosystem because there would be no construction. For the aquatic ecosystem, there would be no opportunities for fish habitat improvement, and continued erosive flows would continue to degrade the habitat quality.

7.2.5 Preferred Alternative

Overall, Dundas AS1 is the preferred alternative. This alternative will have similar impacts on the terrestrial ecosystem as Dundas AS2 and AS3. All three will provide similar terrestrial connectivity improvements through the widened bridge span. Dundas AS1 and AS3 will require the most tree removals for the floodplain improvements and bridge replacement.

8 RECOMMENDATIONS AND PERMITTING

Given the potential and confirmed natural heritage features and functions within the study area, the following recommendations have been identified to reduce long-term impacts to the site:

- A Butternut health assessment should be completed to identify whether the Butternut in question is a hybrid or a pure specimen.
- If the Butternut is a pure specimen, no construction works are to occur within 25 m of the Butternut. Any construction activities occurring within 25 m of the Butternut which could pose harm will be subject to an MECP Notice of Activity to register the project activities.
- Bat acoustic surveys should be completed to confirm the presence or absence of potential SAR bats if an individual tree or forested area identified as potential maternity roosting habitat will be impacted or removed. If SAR bats are present, approval for SAR bat habitat removal from the MECP will be required.
- No tree removals should be performed during the breeding bird window (April 1 to August 30) or the bat roosting window (April 1 to September 30). Should any vegetation clearing be required within the nesting season, a qualified avian biologist shall undertake nest sweeps prior to tree removal to ensure that no nests are present.
- No in-water works should be performed during the fisheries timing window for warm water species (April 1 to June 30). A fish salvage should be conducted prior to the commencement of in-water works, and fish exclusion measures should be installed and maintained throughout the duration of the works.
- Any in-water works will require a Request for Review to the DFO to determine if the project is in contravention of the Fisheries Act.
- Any works within the regulation limit will require a permit through the TRCA.
- Soil best management practices should be implemented during construction to protect the health of the soil and maximize the success of the restoration. This should include an erosion and sediment control plan, proper storage of topsoil for reuse, compaction avoidance measures, topsoil organic testing (if necessary), and topsoil amendments (if necessary).

- Wildlife exclusion fencing should be installed and maintained to ensure no SAR reptiles or amphibians enter the construction site.

9 CONCLUSION

Matrix was retained by the City to create a feasibility study for flood mitigation measures in Little Etobicoke Creek. Little Etobicoke Creek is an urban creek system which has been highly impacted by flood flows in recent years.

This NHS report has been developed in support of the feasibility study to identify the existing conditions of the natural heritage features within the study area. The results of the NHS indicate the presence of the following natural heritage features within the study area:

- Significant woodland
- Significant valleyland (Little Etobicoke Creek)
- Fish and fish habitat
- Confirmed SAR and SCC:
 - ✦ Butternut (SAR)
 - ✦ Chimney Swift (SAR)
 - ✦ Eastern Wood-pewee (SCC)
- Moderate probability of occurrence for six SAR and four SCC

Following the selection of the preferred alternative, a full impact assessment can be completed. The impact assessment will include recommended mitigation and compensation measures to maintain and/or improve the natural heritage features and functions of the study area.

10 REFERENCES

- Brouillet L. et al. 2020. 'Database of Vascular Plants of Canada (VASCAN)'. Online at <http://data.canadensys.net/vascan> and <http://www.gbif.org/dataset/3f8a1297-3259-4700-91fc-acc4170b27ce>, released on 2010-12-10. 2020.
- City of Mississauga. 2020. Mississauga Natural Areas Survey. https://www6.mississauga.ca/onlinemaps/planbldg/nas/key_map/nasmap.pdf
- City of Mississauga. 2023. Mississauga Official Plan. Mississauga, Ontario. 27 July, 2023. <https://www.mississauga.ca/projects-and-strategies/strategies-and-plans/mississauga-official-plan/>

- Dobbyn J.S. 1994. Atlas of the Mammals of Ontario. Federation of Ontario Naturalists. ISBN: 1-896059-02-3. Don Mills, Ontario. 1994. 120 pp.
- Fisheries and Oceans Canada (DFO). 2021. Projects Near Water. Last modified April 14, 2021. <https://www.dfo-mpo.gc.ca/pnw-ppe/index-eng.html>
- Fisheries and Oceans Canada (DFO). 2019. Aquatic species at risk map. Last modified August 23, 2019. <http://www.dfo-mpo.gc.ca/species-especes/sara-lep/map-carte/index-eng.html>
- Important Bird Areas Canada (IBA). 2024. Important Bird and Biodiversity Areas in Canada. Accessed 2024. <https://www.ibacanada.com/>
- Lee H. 2008. Southern Ontario Ecological Land Classification, Vegetation Type List. Prepared for the Ontario Ministry of Natural Resources. London, Ontario. May 2008.
- Lee H. et al. 1998. Ecological Land Classification for Southern Ontario: First Approximation and Its Application. Ontario Ministry of Natural Resources, Southcentral Science Section, Science Development and Transfer Branch. SCSS Field Guide FG-02. 1998.
- Newmaster S.G. et al. 1998. Ontario Plant List. Ontario Forest Research Institute. ISBN: 0-7778-7318-4. Sault Ste. Marie, Ontario. 1998. 550 pp.
- Oldham M.J. et al. 1995. Floristic Quality Assessment System for Southern Ontario. Ontario Ministry of Natural Resources. Natural Heritage Information Centre. Queen's Printer for Ontario. 69 p. 1995.
- Ontario Breeding Bird Atlas (OBBA). 2001. Ontario Breeding Bird Atlas Guide for Participants. Bird Studies Canada, Environment Canada, Federation of Ontario Naturalists, Ministry of Natural Resources, Ontario Field Ornithologists. March 2001.
- Ontario Ministry of Municipal Affairs and Housing (MMAH). 2020. Provincial Policy Statement, 2020. Issued under Section 3 of the Planning Act. Queen's Printer for Ontario, 2020. Toronto, Ontario. 1 May 2020.
- Ontario Ministry of Natural Resources and Forestry (MNRF). 2021. Land Information Ontario. Updated November 22, 2021. Peterborough, Ontario. 2021. <https://www.ontario.ca/page/land-information-ontario>
- Ontario Ministry of Natural Resources and Forestry (MNRF). 2024. Make a Map: Natural Heritage Areas. Mapping application. Accessed March 2024. <http://www.gisoeapp.lrc.gov.on.ca/web/MNR/NHLUPS/NaturalHeritage/Viewer/Viewer.html>

- Ontario Ministry of Natural Resources and Forestry (MNRF). 2017. Survey Protocol for Species at Risk Bats within Treed Habitats Little Brown Myotis, Northern Myotis & Tri-Coloured Bat. Guelph District. April 2017.
- Ontario Ministry of Natural Resources (MNR). 2011. Bat and Bat Habitats: Guidelines for Wind Power Projects. Ontario, Canada. July 2011.
- Ontario Ministry of Natural Resources (MNR). 2010. Natural Heritage Reference Manual for Natural Heritage Policies of the Provincial Policy Statement, 2005. Second Edition. Queen's Printer. Toronto, Ontario. March 18, 2010. 2010.
- Ontario Nature. 2024. Ontario Reptile and Amphibian Atlas. Accessed March 2024.
<https://ontarionature.org/programs/citizen-science/reptile-amphibian-atlas/species/>
- Stanfield L. 2017. Ontario Stream Assessment Protocol (OSAP). Version 10.0. Fisheries Policy Section. Ontario Ministry of Natural Resources. April 2017.
- Toronto and Region Conservation Authority (TRCA). 2019a. Vegetation Communities for the TRCA Jurisdictions. 2019.
- Toronto and Region Conservation Authority (TRCA). 2019b. TRCA Open Data & Information. Accessed July 2019. <https://data.trca.ca/>
- Toronto and Region Conservation Authority (TRCA). 2018. Etobicoke Creek: Watershed Report Card 2018. Downsview, Ontario. 2018. reportcard.trca.ca
- Toronto and Region Conservation Authority (TRCA). 2017. Annual Local Occurrence Score and Local Rank Update: Terrestrial Fauna and Flora Species and Vegetation Communities. Toronto, Ontario. July 2017.
- Toronto and Region Conservation Authority (TRCA). 2010. Etobicoke and Mimico Creek Watersheds, Technical Update Report. 2010.
- Toronto and Region Conservation Authority (TRCA). 2006. GTA Living City Project, Etobicoke Creek, The Aquatic Ecosystem. Downsview, Ontario. 2006.
- Toronto Entomologists' Association (TEA). 2019. Ontario Butterfly Atlas. Updated April 2019. 2019.
http://www.ontarioinsects.org/atlas_online.htm

APPENDIX A
Wildlife Lists

TABLE 1 Bird Species

Species		Conservation Rank					Source						
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC ⁽¹⁾	OBBA ⁽²⁾	eBird ⁽³⁾	iNaturalist ⁽⁴⁾	MECP Information Request (2021)	TRCA Data Request	Matrix Field Observations
Accipitridae Hawks, Kites, Eagles & Allies													
<i>Accipiter cooperii</i>	Cooper's Hawk	S4						x	x	x			
<i>Accipiter striatus</i>	Sharp-shinned Hawk	S5						x	x				
<i>Aquila chrysaetos</i>	Golden Eagle	S1B, S4N	END						x				
<i>Buteo jamaicensis</i>	Red-tailed Hawk	S5						x	x	x			
<i>Buteo platypterus</i>	Broad-winged Hawk	S5B							x				
<i>Circus cyaneus</i>	Northern Harrier	S5B, S4N							x				
<i>Haliaeetus leucocephalus</i>	Bald Eagle	S4	SC					x	x				
Alaudidae Larks													
<i>Eremophila alpestris</i>	Horned Lark	S4						x					
Alcedinidae Kingfishers													
<i>Megaceryle alcyon</i>	Belted Kingfisher	S5B, S4N						x	x	x			x
Apodidae Swifts													
<i>Chaetura pelagica</i>	Chimney Swift	S3B	THR	THR	THR		x	x	x		x	x	x
Anatidae Ducks, Geese & Swans													
<i>Aix sponsa</i>	Wood Duck	S5B, S3N						x	x				
<i>Anas discors</i>	Blue-winged Teal	S3B,S4M						x					
<i>Anas platyrhynchos</i>	Mallard	S5						x	x	x		x	x
<i>Anas rubripes</i>	American Black Duck	S4						x					
<i>Anas strepera</i>	Gadwall	S4B, S4N, S5M						x					
<i>Branta canadensis</i>	Canada Goose	S5						x	x				
<i>Cygnus buccinator</i>	Trumpeter Swan	S4						x					
<i>Cygnus olor</i>	Mute Swan	SNA						x	x				
<i>Lophodytes cucullatus</i>	Hooded Merganser	S5						x					
<i>Melanitta perspicillata</i>	Surf Scoter	S4B/S5N							x				
<i>Cygnus columbianus</i>	Tundra Swan	S2B, S4N, S3M							x				
<i>Mergus merganser</i>	Common Merganser	S5						x	x	x			
Ardeidae Herons and Bitterns													
<i>Ardea herodias</i>	Great Blue Heron	S4							x				
<i>Butorides virescens</i>	Green Heron	S4B						x					
<i>Ixobrychus exilis</i>	Least Bittern	S4B	THR	THR	THR			x					
<i>Nycticorax nycticorax</i>	Black-crowned Night-heron	S3B, S2N, S4M								x			
Bombycillidae Waxwings													
<i>Bombycilla cedrorum</i>	Cedar Waxwing	S5						x	x	x			
Caprimulgidae Nightjars													
<i>Chordeiles minor</i>	Common Nighthawk	S4B	SC	SC	SC			x					
<i>Antrastomus vociferus</i>	Eastern Whip-poor-will	S4B	THR	SC	THR		x						
Cardinalidae Cardinals, Grosbeaks & Allies													
<i>Cardinalis cardinalis</i>	Northern Cardinal	S5						x	x	x		x	x
<i>Passerina cyanea</i>	Indigo Bunting	S5B						x	x	x		x	
<i>Pheucticus ludovicianus</i>	Rose-breasted Grosbeak	S5B						x	x	x			
<i>Piranga olivacea</i>	Scarlet Tanager	S5B						x	x				
Cathartidae Vultures													
<i>Cathartes aura</i>	Turkey Vulture	S5B, S3N						x	x				
Certhiidae Creepers													
<i>Certhia americana</i>	Brown Creeper	S5						x	x	x			x
Charadriidae Plovers													
<i>Charadrius semipalmatus</i>	Semipalmated Plover	S4B, S5M							x				
<i>Charadrius vociferus</i>	Killdeer	S4B						x	x	x			
Columbidae Pigeons & Doves													
<i>Columba livia</i>	Rock Pigeon	SNA						x	x	x			

Species		Conservation Rank					Source						
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC ⁽¹⁾	OBBA ⁽²⁾	eBird ⁽³⁾	iNaturalist ⁽⁴⁾	MECP Information Request (2021)	TRCA Data Request	Matrix Field Observations
<i>Zenaidura macroura</i>	Mourning Dove	S5						x	x	x			
Corvidae													
<i>Corvus brachyrhynchos</i>	Crows & Jays							x	x			x	
<i>Corvus corax</i>	American Crow	S5						x	x				
<i>Corvus ossifragus</i>	Common Raven	S5						x	x				
<i>Corvus ossifragus</i>	Fish Crow	S1B, S3N						x					
<i>Cyanocitta cristata</i>	Blue Jay	S5						x	x			x	x
Cuculidae													
<i>Coccyzus americanus</i>	Cuckoo & Anis												
<i>Coccyzus americanus</i>	Yellow-billed Cuckoo	S4B						x					
<i>Coccyzus erythrophthalmus</i>	Black-billed Cuckoo	S4S5B						x	x				
Emberizidae													
<i>Ammodramus henslowii</i>	New World Sparrows & Allies												
<i>Ammodramus henslowii</i>	Henslow's Sparrow	S1B	END	END	END		x						
<i>Junco hyemalis</i>	Dark-eyed Junco	S5							x	x			
<i>Melospiza georgiana</i>	Swamp Sparrow	S5B, S4N						x	x				
<i>Melospiza lincolni</i>	Lincoln's Sparrow	S5B							x	x			
<i>Melospiza melodia</i>	Song Sparrow	S5						x	x	x			x
<i>Passerculus sandwichensis</i>	Savannah Sparrow	S5B, S3N						x	x				
<i>Passerella iliaca</i>	Fox Sparrow	S5B, S3N							x				
<i>Pipilo erythrophthalmus</i>	Eastern Towhee	S4B, S3N						x	x				
<i>Spizella arborea</i>	American Tree Sparrow	S5							x				
<i>Spizella pallida</i>	Clay-colored Sparrow	S4B							x				
<i>Spizella passerina</i>	Chipping Sparrow	S5B, S3N						x	x	x		x	x
<i>Zonotrichia albicollis</i>	White-throated Sparrow	S5							x	x			
<i>Zonotrichia leucophrys</i>	White-crowned Sparrow	S5B, S3N							x				
<i>Spizella pusilla</i>	Field Sparrow	S4B, S3N						x	x				
Falconidae													
<i>Falco columbarius</i>	Carcaras & Falcons												
<i>Falco columbarius</i>	Merlin	S5						x	x	x			
<i>Falco peregrinus</i>	Peregrine Falcon	S4	SC					x		x	x		
<i>Falco sparverius</i>	American Kestrel	S4						x	x				
Fringillidae													
<i>Acanthis flammea</i>	Finches & Allies												
<i>Acanthis flammea</i>	Common Redpoll	S5							x				
<i>Haemorhous mexicanus</i>	House Finch	SNA						x	x	x			
<i>Haemorhous purpureus</i>	Purple Finch	S5							x				
<i>Spinus pinus</i>	Pine Siskin	S5							x				
<i>Spinus tritis</i>	American Goldfinch	S5						x	x			x	x
Gaviidae													
<i>Gavia immer</i>	Loons												
<i>Gavia immer</i>	Common Loon	S5							x				
Gruidae													
<i>Grus canadensis</i>	Cranes												
<i>Grus canadensis</i>	Sandhill Crane	S5B, S3N							x				
Hirundinidae													
<i>Hirundo rustica</i>	Swallows												
<i>Hirundo rustica</i>	Barn Swallow	S4B	SC	SC	THR		x	x	x		x		
<i>Petrochelidon pyrrhonota</i>	Cliff Swallow	S4S5B						x					
<i>Progne subis</i>	Purple Martin	S3B						x	x				
<i>Riparia riparia</i>	Bank Swallow	S4B	THR	THR	THR			x			x		
<i>Stelgidopteryx serripennis</i>	Northern Rough-winged Swallow	S4B						x	x				
<i>Tachycineta bicolor</i>	Tree Swallow	S4S5B						x	x				
Icteridae													
<i>Agelaius phoeniceus</i>	New World Blackbird												
<i>Agelaius phoeniceus</i>	Red-winged Blackbird	S5						x	x	x		x	x
<i>Dolichonyx oryzivorus</i>	Bobolink	S4B	THR	SC	THR			x					
<i>Euphagus carolinus</i>	Rusty Blackbird	S4B, S3N	SC	SC	SC				x				
<i>Icterus galbula</i>	Baltimore Oriole	S4B						x	x	x		x	
<i>Icterus spurius</i>	Orchard Oriole	S4B						x	x				
<i>Molothrus ater</i>	Brown-headed Cowbird	S5						x	x			x	
<i>Quiscalus quiscula</i>	Common Grackle	S5						x	x	x		x	x
<i>Sturnella magna</i>	Eastern Meadowlark	S4B, S3N	THR	THR	THR			x					

Species		Conservation Rank					Source						
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC ⁽¹⁾	OBBA ⁽²⁾	eBird ⁽³⁾	iNaturalist ⁽⁴⁾	MECP Information Request (2021)	TRCA Data Request	Matrix Field Observations
Laridae													
Gulls, Terns & Skimmers													
<i>Larus argentatus</i>	Herring Gull	S4B, S5N							x				
<i>Larus delawarensis</i>	Ring-billed Gull	S5							x	x			
<i>Sterna hirundo</i>	Common Tern	S4B						x					
Mimidae													
Mockingbirds, Thrashers & Allies													
<i>Dumetella carolinensis</i>	Gray Catbird	S5B, S3N						x	x			x	x
<i>Mimus polyglottos</i>	Northern Mockingbird	S4						x	x	x			
<i>Toxostoma rufum</i>	Brown Thrasher	S4B						x	x				
Motacillidae													
<i>Anthus rubescens</i>	American Pipit	S4B							x				
Pandionidae													
Osprey													
<i>Pandion haliaetus</i>	Osprey	S5B						x					
Paridae													
Chickadees and Titmice													
<i>Poecile atricapillus</i>	Black-capped Chickadee	S5						x	x	x		x	x
Parulidae													
Wood Warblers													
<i>Cardellina canadensis</i>	Canada Warbler	S5B	SC	SC	THR				x		x		
<i>Cardellina pusilla</i>	Wilson's Warbler	S5B							x				
<i>Geothlypis philadelphia</i>	Mourning Warbler	S5B						x	x				
<i>Geothlypis trichas</i>	Common Yellowthroat	S5B, S3N						x	x			x	
<i>Mniotilta varia</i>	Black-and-white Warbler	S5B						x	x	x			x
<i>Oreothlypis celata</i>	Orange-crowned Warbler	S5B							x				
<i>Oreothlypis peregrina</i>	Tennessee Warbler	S5B							x				
<i>Oreothlypis ruficapilla</i>	Nashville Warbler	S5B							x				
<i>Seiurus aurocapilla</i>	Ovenbird	S5B						x	x				
<i>Setophaga americana</i>	Northern Parula	S5B							x	x			
<i>Setophaga caerulescens</i>	Black-throated Blue Warbler	S5B						x	x	x			
<i>Setophaga castanea</i>	Bay-breasted Warbler	S5B							x				
<i>Setophaga coronata</i>	Yellow Rumped Warbler	S5B, S4N							x	x			
<i>Setophaga fusca</i>	Blackburnian Warbler	S5B						x	x	x			
<i>Setophaga magnolia</i>	Magnolia Warbler	S5B							x	x			
<i>Setophaga pennsylvanica</i>	Chestnut-sided warbler	S5B						x	x				
<i>Setophaga petechai</i>	Yellow Warbler	S5B						x	x				x
<i>Setophaga pinus</i>	Pine Warbler	S5B, S3N						x	x				
<i>Setophaga ruticilla</i>	American Redstart	S5B						x	x				
<i>Setophaga striata</i>	Blackpoll Warbler	S5B							x				
<i>Setophaga tigrina</i>	Cape May Warbler	S5B							x	x			
<i>Setophaga virens</i>	Black-throated Green Warbler	S5B							x	x			
Passeridae													
Sparrows													
<i>Passer domesticus</i>	House Sparrow	SNA							x	x	x	x	
Phalacrocoracidae													
Cormorants													
<i>Phalacrocorax auritus</i>	Double-crested Cormorant	S5B, S4N							x	x			
Phasianidae													
Partridges, Grouse, Turkeys													
<i>Meleagris gallopavo</i>	Wild Turkey	S5							x				
Picidae													
Woodpeckers													
<i>Colaptes auratus</i>	Northern Flicker	S5							x	x	x		
<i>Dryocopus pileatus</i>	Pileated Woodpecker	S5								x			
<i>Leuconotopicus villosus</i>	Hairy Woodpecker	S5							x	x		x	x
<i>Melanerpes carolinus</i>	Red-bellied Woodpecker	S5							x	x			
<i>Melanerpes erythrocephalus</i>	Red-headed Woodpecker	S3	END	END	END		x						
<i>Picoides pubescens</i>	Downy Woodpecker	S5							x	x			x
<i>Sphyrapicus varius</i>	Yellow-bellied Sapsucker	S5B, S3N							x	x			x
Podicipedidae													
Grebes													
<i>Podiceps grisegena</i>	Red-necked Grebe	S3							x				
<i>Podilymbus podiceps</i>	Pied-billed Grebe	S4B, S2N							x				

Species		Conservation Rank					Source						
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC ⁽¹⁾	OBBA ⁽²⁾	eBird ⁽³⁾	iNaturalist ⁽⁴⁾	MECP Information Request (2021)	TRCA Data Request	Matrix Field Observations
Poliptilidae													
Gnatcatchers													
<i>Poliptila caerulea</i>	Blue-gray Gnatcatcher	S4B						x	x				
Rallidae													
Railes, Gallinules & Coots													
<i>Gallinula chloropus</i>	Common Gallinule	S3B						x					
<i>Rallus limicola</i>	Virginia Rail	S4S5B						x					
Regulidae													
Kinglets													
<i>Regulus calendula</i>	Ruby-crowned Kinglet	S5B, S3N							x				
<i>Regulus satrapa</i>	Golden-crowned Kinglet	S5							x	x			x
Scolopacidae													
Sandpipers, Phalaropes & Allies													
<i>Actitis macularius</i>	Spotted Sandpiper	S5B						x		x			
<i>Scolopax minor</i>	American Woodcock	S4B						x		x			
<i>Tringa solitaria</i>	Solitary Sandpiper	S4B, S5M								x			
Sittidae													
Nuthatches													
<i>Sitta canadensis</i>	Red-breasted Nuthatch	S5						x	x	x			
<i>Sitta carolinensis</i>	White-breasted Nuthatch	S5						x	x				x
Stercorariidae													
Skuas													
<i>Bubo virginianus</i>	Great Horned Owl	S4						x	x	x			
<i>Megascops asio</i>	Eastern Screech Owl	S4								x			
Sturnidae													
Starlings													
<i>Sturnus vulgaris</i>	European Starling	SNA						x	x	x			
Trochillidae													
Hummingbirds													
<i>Archilochus colubris</i>	Ruby-throated Hummingbird	S5B						x	x				
Troglodytidae													
Wrens													
<i>Cistothorus palustris</i>	Marsh Wren	S4B, S3N						x					
<i>Thyrothorus ludovicianus</i>	Carolina Wren	S4						x	x				
<i>Troglodytes aedon</i>	House Wren	S5B						x	x				
<i>Troglodytes hiemalis</i>	Winter Wren	S5B, S4N							x				
Turdidae													
Thrushes													
<i>Catharus fuscescens</i>	Veery	S5B						x	x	x			
<i>Catharus guttatus</i>	Hermit Thrush	S5B, S4N							x	x			
<i>Catharus minimus</i>	Gray-cheeked Thrush	S4?B, S4M							x				
<i>Catharus ustulatus</i>	Swainson's Thrush	S5B							x	x			
<i>Hyllocichla mustelina</i>	Wood Thrush	S4B	SC	THR	THR			x	x		x		
<i>Sialia sialis</i>	Eastern Bluebird	S5B, S4N						x					
<i>Turdus migratorius</i>	American Robin	S5						x	x	x		x	x
Tyrannidae													
Tyrant Flycatchers													
<i>Contopus virens</i>	Eastern Wood-pewee	S4B	SC	SC	SC		x	x	x		x		x
<i>Empidonax alnorum</i>	Alder Flycatcher	S5B							x				
<i>Empidonax flaviventris</i>	Yellow-bellied Flycatcher	S5B							x				
<i>Empidonax minimus</i>	Least Flycatcher	S5B						x	x				x
<i>Empidonax traillii</i>	Willow Flycatcher	S4B						x	x				
<i>Myiarchus crinitus</i>	Great Crested Flycatcher	S5B						x	x				x
<i>Sayornis phoebe</i>	Eastern Phoebe	S5B						x	x				
<i>Tyrannus tyrannus</i>	Eastern Kingbird	S4B						x	x	x			
Vireonidae													
Vireos													
<i>Vireo gilvus</i>	Warbling Vireo	S5B						x	x				
<i>Vireo olivaceus</i>	Red-eyed Vireo	S5B						x	x			x	
<i>Vireo solitarius</i>	Blue-headed Vireo	S5B							x	x			
<i>Vireo flavifrons</i>	Yellow-throated Vireo	S4B						x					
<i>Vireo philadelphicus</i>	Philadelphia Vireo	S5B							x				
Total							6	110	131	53	7	19	24

Notes:

S-rank

S1 - Critically Imperiled

COSEWIC

NAR - Not at Risk

Species		Conservation Rank					Source						
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC ⁽¹⁾	OBBA ⁽²⁾	eBird ⁽³⁾	iNaturalist ⁽⁴⁾	MECP Information Request (2021)	TRCA Data Request	Matrix Field Observations
S2 - Imperiled	SC - Special Concern												
S3 - Vulnerable	THR - Threatened												
S4 - Apparently Secure	END - Endangered												
S5 - Secure	EXT - Extinct												
SU - Unrankable	EXP - Extirpated												
SNA - Unranked	DD - Data Deficient												
SX - Presumed Extirpated													
SH - Possibly Extirpated	<u>SARA Schedule</u>												
S#? - Rank Uncertain	Schedule 1 - Officially protected under SARA												
	Schedule 2 - threatened/endangered; may be reassessed for consideration for inclusion to Schedule 1												
	Schedule 3 - special concern; may be reassessed for consideration for inclusion to Schedule 1												
<u>ESA</u>													
SC - Special Concern													
THR - Threatened	<u>COSSARO</u>												
END - Endangered	NAR - Not at Risk												
EXT - Extinct	SC - Special Concern												
EXP - Extirpated	THR - Threatened												
	END - Endangered												
	EXP - Extirpated												
	DD - Data Deficient												
<u>Additional Notes</u>													
ESA - <i>Endangered Species Act</i>													
COSEWIC - Committee on the Status of Endangered Wildlife in Canada													
SARA - <i>Species at Risk Act</i>													
NHIC - Natural Heritage Information Centre													
OBBA - Ontario Breeding Bird Atlas													

Sources:

1 Ontario Ministry of Natural Resources and Forestry (MNRF). 2024. *Make a Map: Natural Heritage Areas*. Mapping application. Accessed March 2024. https://www.lioapplications.lrc.gov.on.ca/Natural_Heritage/index.html?viewer=Natural_Heritage.Natural_Heritage&locale=en-CA

2 Birds Canada et al. 2024. *Ontario Breeding Bird Atlas Data Summary Tool*. NatureCounts platform. Accessed March 2024. <https://naturecounts.ca/nc/onatlas/findsquare.jsp>

3 Cornell Lab of Ornithology. 2024. *eBird*. Accessed March 2024. <https://ebird.org/home>

3 iNaturalist. 2024. *Observations*. Accessed March 2024. <https://www.inaturalist.org/observations>

TABLE 2 Reptile and Amphibian Species

Species		Conservation Rank					Source			
Scientific Name	Common Name	Provincial (S-RANK)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Regional (Peel)	NHIC	ORAA	iNaturalist	Matrix Observations
Cryptodeira		Turtles								
<i>Chelydra serpentina</i>	Snapping Turtle	S4	SC	SC	SC		x	x		
<i>Chrysemys picta marginata</i>	Midland Painted Turtle	S4		SC	SC		x	x	x	
<i>Emydoidea blandingii</i>	Blanding's Turtle	S3	THR	END	END			x		
<i>Graptemys geographica</i>	Northern Map Turtle	S3	SC	SC	SC			x	x	
<i>Sternotherus odoratus</i>	Eastern Musk Turtle	S3	SC	SC	SC			x		
<i>Trachemys scripta elegans</i>	Red-eared Slider	SNA						x		
Squamata		Snakes								
<i>Diadophis punctatus</i>	Ring-necked Snake	S4						x		
<i>Lampropeltis triangulum</i>	Eastern Milksnake	S4		SC	SC		x	x	x	
<i>Nerodia sipedon sipedon</i>	Northern Watersnake	S5						x	x	
<i>Ophedrys vernalis</i>	Smooth Greensnake	S4						x		
<i>Storeria dekayi</i>	DeKay's Brownsnake	S5						x	x	
<i>Storeria occipitomaculata</i>	Red-bellied snake	S5						x		
<i>Thamnophis sirtalis sirtalis</i>	Eastern Gartersnake	S5						x	x	
Caudata		Salamanders								
<i>Ambystoma hybrid pop. 1</i>	Jefferson X Blue-spotted Salamander, Jefferson genome dominates	S2	END	END	END			x		
<i>Ambystoma jeffersonianum</i>	Jefferson Salamander	S2	END	END	END			x		
<i>Ambystoma maculatum</i>	Spotted Salamander	S4						x		
<i>Necturus maculosus</i>	Mudpuppy	S4						x		
<i>Notophthalmus viridescens viridescens</i>	Red-spotted Newt	S5						x		
<i>Plethodon cinereus</i>	Eastern Red-backed Salamander	S5						x	x	
Anura		Frogs and Toads								
<i>Anaxyrus americanus</i>	American Toad	S5						x	x	
<i>Hyla versicolor</i>	Gray Treefrog	S5						x		
<i>Lithobates catesbeianus</i>	American Bullfrog	S4						x		
<i>Lithobates clamitans</i>	Green Frog	S5						x		
<i>Lithobates palustris</i>	Pickerel Frog	S4						x		
<i>Lithobates pipiens</i>	Northern Leopard Frog	S5						x		
<i>Lithobates sylvaticus</i>	Wood Frog	S5						x		
<i>Pseudacris crucifer</i>	Spring Peeper	S5						x		
<i>Pseudacris trisetaria pop. 1</i>	Western Chorus Frog (Great Lakes / St. Lawrence - Canadian Shield population)	S4		THR	THR			x		
Total:							3	28	8	0

TABLE 3 Fish Species

Scientific Name	Species Name	Common Name	Conservation Rank					Source				
			Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC (1)	iNaturalist (2)	DFO (3)	LIO (4)	Etobicoke Creek Fish Data (TRCA)
Clupeiformes												
<i>Alosa pseudoharengus</i>	Alewife		SNA									x
<i>Dorosoma cepedianum</i>	Gizzard Shad		S4									x
Cypriniformes												
<i>Campostoma anomalum</i>	Central Stoneroller		S4									x
<i>Carassius auratus</i>	Goldfish		SNA						x			x
<i>Caesius plumbeus</i>	Lake Chub		S5									x
<i>Cyprinus carpio</i>	Common Carp		SNA									x
<i>Natemitogon crysoleucas</i>	Golden Shiner		S5								x	
<i>Natrapis atherinoides</i>	Emerald Shiner		S5								x	
<i>Natrapis heterodon</i>	Blackchin Shiner		S4									x
<i>Natrapis heterolepis</i>	Blacknose Shiner		S5									x
<i>Natrapis hudsonius</i>	Spottail Shiner		S5									x
<i>Natrapis stramineus</i>	Sand Shiner		S4									x
<i>Chrasomus eos</i>	Northern Redbelly Dace		S5									x
<i>Chrasomus neogaeus</i>	Finescale Dace		S5						x			
<i>Pimephales notatus</i>	Bluntnose Minnow		S5						x	x		x
<i>Pimephales promelas</i>	Fathead Minnow		S5						x			x
<i>Rhinichthys atratulus</i>	Western Blacknose Dace		S5						x	x		x
<i>Rhinichthys cataractae</i>	Longnose Dace		S5						x	x		x
<i>Semotilus atromaculatus</i>	Creek Chub		S5						x	x		x
<i>Luxilus cornutus</i>	Common Shiner		S5						x	x		x
<i>Margariscus margarita</i>	Northern Pearl Dace		S5									x
<i>Catostomus commersonii</i>	White Sucker		S5						x	x		x
<i>Hypentelium nigricans</i>	Northern Hog Sucker		S4									x
Esociformes												
<i>Umbra limi</i>	Central Mudminnow		S5									x
<i>Esox lucius</i>	Northern Pike		S5									x
Gasterosteiformes												
<i>Culaea inconstans</i>	Brook Stickleback		S5							x		x
<i>Gasterosteus aculeatus</i>	Threespine Stickleback		S4S5							x		x
Perciformes												
<i>Aplodinotus grunniens</i>	Freshwater Drum		S5									x
<i>Morone americana</i>	White Perch		SNA									x
<i>Ambloplites rupestris</i>	Rock Bass		S5							x		x
<i>Lepomis gibbosus</i>	Pumpkinseed		S5							x		x
<i>Micropterus dolomieu</i>	Smallmouth Bass		S5									x
<i>Micropterus salmoides</i>	Largemouth Bass		S5									x
<i>Pomoxis nigromaculatus</i>	Black Crappie		S4									x
<i>Etheostoma flabellare</i>	Fantail Darter		S4									x
<i>Etheostoma nigrum</i>	Johnny Darter		S5									x
<i>Etheostoma olmstedii</i>	Tessellated Darter		S4									x
<i>Perca flavescens</i>	Yellow Perch		S5							x		
<i>Neogobius melanostomus</i>	Round Goby		SNA									x
Percopsiformes												
<i>Percopsis omiscomaycus</i>	Trout-perch		S5									x
Salmoniformes												
<i>Oncorhynchus tshawytscha</i>	Chinook Salmon		SNA									x
<i>Oncorhynchus mykiss</i>	Rainbow Trout		SNA									x
<i>Salmo trutta</i>	Brown Trout		S5									x
Scorpaeniformes												
<i>Cottus bairdii</i>	Mottled Sculpin		S5									x
Siluriformes												
<i>Ameiurus nebulosus</i>	Brown Bullhead		S5									x
Anguilliformes												
<i>Anguilla rostrata</i>	American Eel		S1S2	END	THR							
<i>Clinostomus elongatus</i>	Redside Dace		S1	END	END	END						
Acipenseriformes												
<i>Acipenser fulvescens</i> pop. 1	Lake Sturgeon (Saskatchewan - Nelson River population)		S2	THR	END							
<i>Acipenser fulvescens</i> pop. 2	Lake Sturgeon (Southern Hudson Bay - James Bay population)		S3	SC	SC	SC						
<i>Acipenser fulvescens</i> pop. 3	Lake Sturgeon (Great Lakes - Upper St. Lawrence River population)		S2	END	THR							
<i>Polyodon spathula</i>	Paddlefish		SX	EXP	EXP	EXP						
Amiiformes												
<i>Amia calva</i>	Bowfin		S4									
Alberrniformes												
<i>Labidesthes sicculus</i>	Brook Silverside		S4									
Characiformes												
<i>Piaractus brachipomus</i>	Redbelly Pacu		SNA									
<i>Ctenopharyngodon idella</i>	Grass Carp		SNA									
<i>Exoglossum maxillingua</i>	Cutlip Minnow		S2	THR	SC	SC						
<i>Hybognathus hankinsoni</i>	Brassy Minnow		S5									
<i>Hybognathus regius</i>	Eastern Silvery Minnow		S2									
<i>Nocomis biguttatus</i>	Hornyhead Chub		S4									
<i>Nocomis micropogon</i>	River Chub		S4									
<i>Natropis anogenus</i>	Pugnose Shiner		S2	THR	THR	THR						

Scientific Name	Species Name Common Name	Conservation Rank					Source						
		Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC (1)	iNaturalist (2)	DFO (3)	LO (4)	Etobicoke Creek Fish Data (TRCA)	Etobicoke Management Plan (TRCA)	Matrix Field Observations
<i>Natropis bifrenatus</i>	Bridle Shiner	S2	SC	SC	SC								
<i>Natropis buchonani</i>	Ghost Shiner	S3											
<i>Natropis photogenis</i>	Silver Shiner	S2S3	THR	THR	THR								
<i>Natropis rubellus</i>	Rosyface Shiner	S4											
<i>Natropis voluceilus</i>	Mimic Shiner	S5											
<i>Scardinus erythrophthalmus</i>	Rudd	SNA											
<i>Semotilus corporalis</i>	Fallfish	S4											
<i>Cyprinella spiloptera</i>	Spotfin Shiner	S4											
<i>Erimystax punctatus</i>	Gravel Chub	SX	EXP	EXP	EXP								
<i>Luxilus chrysocephalus</i>	Striped Shiner	S4											
<i>Lythrurus umbratilis</i>	Redfin Shiner	S4											
<i>Macrhybopsis storeriana</i>	Silver Chub (Great Lakes - Upper St. Lawrence populations)	S2	THR	END	END								
<i>Osgoodeus emiliae</i>	Pugnose Minnow	S2	THR	THR	THR								
<i>Carpodes cyprinus</i>	Quillback	S4											
<i>Catostomus catostomus</i>	Longnose Sucker	S5											
<i>Erimyzon sucetta</i>	Lake Chubsucker	S2	END	END	END								
<i>Ictiobus cyprinellus</i>	Bigmouth Buffalo	S27											
<i>Ictiobus niger</i>	Black Buffalo	SNA			SC								
<i>Minytrema melanops</i>	Spotted Sucker	S2	SC	SC	SC								
<i>Maxostoma anisurum</i>	Silver Redhorse	S4											
<i>Maxostoma carinatum</i>	River Redhorse	S2	SC	SC	SC								
<i>Maxostoma duquesnei</i>	Black Redhorse	S2	THR	THR	THR								
<i>Maxostoma erythrurum</i>	Golden Redhorse	S4											
<i>Maxostoma macrolepidotum</i>	Shorthead Redhorse	S5											
<i>Maxostoma valenciennesi</i>	Greater Redhorse	S3											
<i>Natropis biennis</i>	River Shiner	S1?											
Cyprinodontiformes													
<i>Fundulus diaphanus</i>	Banded Killifish	S5											
<i>Fundulus notatus</i>	Blackstripe Topminnow	S2	SC	SC	SC								
<i>Gambusia affinis</i>	Western Mosquitofish	SNA											
<i>Esox americanus vermiculatus</i>	Grass Pickerel	S3	SC	SC	SC								
<i>Dallia pectoralis</i>	Alaska Blackfish	SNA											
<i>Esox masquinongy</i>	Muskellunge	S4											
<i>Esox niger</i>	Chain Pickerel	SNA											
Gadiformes													
<i>Lota lota</i>	Burbot	S5											
<i>Apeltes quadracus</i>	Fourspine Stickleback	SNA											
<i>Pungitius pungitius</i>	Ninespine Stickleback	S5											
Hiodontiformes													
<i>Hiodon alosoides</i>	Goldeye	S3											
<i>Hiodon tergisus</i>	Mooneye	S4											
Lepisosteiformes													
<i>Lepisosteus oculatus</i>	Spotted Gar	S1	END	END	END								
<i>Lepisosteus osseus</i>	Longnose Gar	S4											
<i>Lepisosteus platyhincus</i>	Florida Gar	SNA											
Osmeriformes													
<i>Osmerus mordax</i>	Rainbow Smelt	S5											
<i>Astronotus ocellatus</i>	Oscar	SNA											
<i>Parachromis managuensis</i>	Jaguar Guapote	SNA											
<i>Morone chrysops</i>	White Bass	S4											
<i>Lepomis cyanellus</i>	Green Sunfish	S4											
<i>Lepomis gulosus</i>	Warmouth	S1	END	END	SC								
<i>Lepomis humilis</i>	Orangespotted Sunfish	SNA											
<i>Lepomis macrochirus</i>	Bluegill	S5											
<i>Pomoxis annularis</i>	White Crappie	S4											
<i>Etheostoma biennioides</i>	Greenside Darter	S4											
<i>Etheostoma caeruleum</i>	Rainbow Darter	S4											
<i>Etheostoma exile</i>	Iowa Darter	S5											
<i>Etheostoma micropetca</i>	Least Darter	S4											
<i>Ammocrypta pellucida</i>	Eastern Sand Darter	S2	END	THR	THR								
<i>Percina caprodes</i>	Loggerhead	S5											
<i>Percina capelandi pop. 1</i>	Channel Darter (Lake Erie population)	S1	SC	END	END								
<i>Percina maculata</i>	Blackside Darter	S4											
<i>Percina shumardi</i>	River Darter	S1	END	END									
<i>Sander canadensis</i>	Sauger	S4											
<i>Sander vitreus vitreus</i>	Walleye	S5											
<i>Sander vitreus glaucus</i>	Blue Pike	SX		EXT									
<i>Gymnocephalus cernuus</i>	Ruffe	SNA											
<i>Proterorhinus marmoratus</i>	Tubenose Goby	SNA											
<i>Lepomis peltastes</i>	Northern Sunfish	S3	SC	SC	SC								
Petromyzontiformes													
<i>Ichthyomyzon fossor</i>	Northern Brook Lamprey	S3	SC	SC	SC								
<i>Lethenteron appendix</i>	American Brook Lamprey	S3											
<i>Petromyzon marinus</i>	Sea Lamprey	SNA											
<i>Ichthyomyzon castaneus pop. 1</i>	Chestnut Lamprey (Great Lakes-St Lawrence population)	S3		DD									
<i>Ichthyomyzon unicuspis pop. 1</i>	Silver Lamprey (Great Lakes - Upper St. Lawrence populations)	S3	SC	SC	SC								

Scientific Name	Species Name Common Name	Conservation Rank					Source						
		Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC (1)	iNaturalist (2)	DFO (3)	LIO (4)	Etobicoke Creek Fish Data (TRCA)	Etobicoke Management Plan (TRCA)	Matrix Field Observations
<i>Ichthyomyzon unicuspis</i> pop. 2	Silver Lamprey (Saskatchewan - Nelson River population)	SU		DD									
<i>Ichthyomyzon castaneus</i> pop. 2	Chestnut Lamprey (Saskatchewan - Nelson River population)	S1?		DD									
Plauronectiformes													
<i>Platichthys flesus</i>	European Flounder	SNA											
<i>Caregonus kiyi</i> <i>kiyi</i>	Upper Great Lakes Kiyi	S3	SC	SC	SC								
<i>Caregonus kiyi orientalis</i>	Lake Ontario Kiyi	SX		EXT									
<i>Salmo salar</i> pop. 2	Atlantic Salmon - Lake Ontario Population	SX		EXT									
<i>Caregonus artedii</i>	Cisco	S5											
<i>Caregonus clupeaformis</i>	Lake Whitefish	S5											
<i>Caregonus hoyi</i>	Bloater	S4											
<i>Caregonus johanna</i>	Deepwater Cisco	SX		EXT									
<i>Caregonus nigripinnis</i>	Blackfin Cisco	SU		DD									
<i>Caregonus reighardi</i>	Shortnose Cisco	SH	END	END	END								
<i>Caregonus zenithicus</i>	Shortjaw Cisco	S2	THR	THR									
<i>Oncorhynchus gorbuscha</i>	Pink Salmon	SNA											
<i>Oncorhynchus keta</i>	Chum Salmon	SNA											
<i>Oncorhynchus kisutch</i>	Coho Salmon	SNA											
<i>Oncorhynchus nerka</i>	Sockeye Salmon	SNA											
<i>Prosopium coulterii</i>	Pygmy Whitefish	SU											
<i>Prosopium cylindraceum</i>	Round Whitefish	S4											
<i>Salvelinus alpinus</i>	Arctic Char	SU											
<i>Salvelinus fontinalis fontinalis</i>	Brook Trout	S5											
<i>Salvelinus namaycush</i>	Lake Trout	S5											
<i>Thymallus arcticus</i>	Arctic Grayling	SNA											
<i>Salvelinus fontinalis timagamiensis</i>	Aurora Trout	S1											
<i>Caregonus clupeaformis</i> pop. 1	Lake Whitefish (Lake Simcoe population)	SNR		DD									
<i>Salvelinus namaycush x fontinalis</i>	Splake	SNA											
<i>Cyclopterus lumpus</i>	Lumpfish	SNA											
<i>Cottus cognatus</i>	Slimy Sculpin	S5											
<i>Cottus ricei</i>	Spoonhead Sculpin	S4											
<i>Myoxocephalus quadricornis</i>	Fourhorn Sculpin	S2?											
<i>Myoxocephalus thompsonii</i>	Deepwater Sculpin (Great Lakes - Western St. Lawrence Populations)	S3?		SC	SC								
<i>Ictalurus punctatus</i>	Channel Catfish	S4											
<i>Naturus flavus</i>	Stonecat	S4											
<i>Naturus gyrinus</i>	Tadpole Madtom	S4											
<i>Naturus insignis</i>	Margined Madtom	SU		DD									
<i>Naturus miurus</i>	Brindled Madtom	S2											
<i>Naturus stigmatosus</i>	Northern Madtom	S1	END	END	END								
<i>Pylodictis olivaris</i>	Flathead Catfish	SNA		DD									
<i>Ameiurus melas</i>	Black Bullhead	S4											
<i>Ameiurus natalis</i>	Yellow Bullhead	S4											
<i>Panaque nigrolineatus</i>	Royal Panaque	SNA											
<i>Alosa sapidissima</i>	American Shad	SNA											
<i>Ictiobus bubalus</i>	Smallmouth Buffalo	SNA											
<i>Percina copelandi</i> pop 2	Channel Darter (Lake Ontario population)	S2	SC	END	END								
<i>Percina copelandi</i> pop 3	Channel Darter (St. Lawrence population)	S3	SC	SC	SC								
Total							0	0	0	15	9	43	6

Notes:

S-rank
 S1 - Critically Imperiled
 S2 - Imperiled
 S3 - Vulnerable
 S4 - Apparently Secure
 S5 - Secure
 SU - Unrankable
 SNA - Unranked
 SX - Presumed Extirpated
 SH - Possibly Extirpated
 SRF - Rank Uncertain

COSEWIC
 NAR - Not at Risk
 SC - Special Concern
 THR - Threatened
 END - Endangered
 EXT - Extinct
 EXP - Extirpated
 DD - Data Deficient

SARA Schedule
 Schedule 1 - Officially protected under SARA
 Schedule 2 - threatened/endangered; may be reassessed for consideration for inclusion to Schedule 1
 Schedule 3 - special concern; may be reassessed for consideration for inclusion to Schedule 1

ESA
 SC - Special Concern
 THR - Threatened
 END - Endangered
 EXT - Extinct
 EXP - Extirpated

COSSARO
 NAR - Not at Risk
 SC - Special Concern
 THR - Threatened
 END - Endangered
 EXP - Extirpated
 DD - Data Deficient

Additional Notes
 ESA - Endangered Species Act
 COSEWIC - Committee on the Status of Endangered Wildlife in Canada
 SARA - Species at Risk Act
 NHIC - Natural Heritage Information Centre
 DFO - Fisheries and Oceans Canada
 LIO - Land Information Ontario

Sources:

Scientific Name	Species Name		Conservation Rank					Source					
	Common Name		Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	Local	NHIC ⁽¹⁾	iNaturalist ⁽²⁾	DFO ⁽³⁾	LIO ⁽⁴⁾	Etobicoke Creek Fish Data (TRCA)	Etobicoke Management Plan (TRCA)

1 Ontario Ministry of Natural Resources and Forestry (MNRF). 2024. Make a Map: Natural Heritage Areas. Mapping application. Accessed March 2024. https://www.lioapplications.lrc.gov.on.ca/Natural_Heritage/index.html?viewer=Natural_Heritage-Natural_Heritage&locale=en-CA

2 iNaturalist. 2024. Observations. Accessed March 2024. <https://www.inaturalist.org/observations>

3 Fisheries and Oceans Canada (DFO). 2024. Aquatic Species at Risk Map. Accessed March 2024. <https://www.dfo-mpo.gc.ca/species-especes/sara-lep/map-carte/index-eng.html>

4 Ontario Ministry of Natural Resources and Forestry (MNRF). 2024. Ontario GeoHub. Open data resource through Land Information Ontario. Accessed March 2024. <https://geohub.lio.gov.on.ca/>

TABLE 4 Insect Species

Species Name		Conservation Rank				Source				
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	NHIC ⁽¹⁾	Ontario Butterfly Atlas ⁽²⁾	iNaturalist ⁽³⁾	Ontario Moth Atlas ⁽⁴⁾	Matrix Field Observations
Coleoptera		Beetles								
<i>Acalymma vittatum</i>	Leaf Beetle	SNR						x		
<i>Carabus granulatus</i>	Ground Beetle	SNA						x		
<i>Cicindela sexguttata</i>	Tiger Beetle	S5						x		
<i>Coccinella septempunctata</i>	Seven-spotted Ladybird Beetle	SNA						x		
<i>Coleomegilla maculata</i>	Spotted Ladybird Beetle	S5						x		
<i>Popillia japonica</i>	Scarab Beetle	SNA						x		
Diptera		Flies								
<i>Xenox tigrinus</i>	Tiger Bee Fly	S3S4						x		
<i>Philaenus spumarius</i>	Meadow Spittlebug	SNA						x		
Hymenoptera		Sawflies, Wasps, Bees, and Ants								
<i>Agapostemon sericeus</i>	Whitish Sweat Bee	S5						x		
<i>Andrena wilkella</i>	European Legume Miner Bee	SNA						x		
<i>Apis mellifera</i>	European Honey Bee	SNA						x		
<i>Bombus bimaculatus</i>	Two-spotted Bumble Bee	S5						x		
<i>Bombus griseocollis</i>	Brown-belted Bumble Bee	S5						x		
<i>Bombus impatiens</i>	Common Eastern Bumble Bee	S5						x		
<i>Megachile sculpturalis</i>	Giant Leafcutter Bee	SNA						x		
<i>Myrmica rubra</i>	European Fire Ant	SNA						x		
<i>Xylocopa virginica</i>	Virginia Carpenter Bee	S4S5						x		
Lepidoptera		Butterflies								
<i>Aglais milberti</i>	Milbert's Tortoiseshell	S5					x			
<i>Amphion floridensis</i>	Nessus Sphinx	S4							x	
<i>Anatrytone logan</i>	Delaware Skipper	S4					x			
<i>Ancyloxypha numitor</i>	Least Skipper	S5					x			
<i>Antheraea polyphemus</i>	Polyphemus Moth	S5						x	x	
<i>Apantesis nais</i>	Nais Tiger Moth	SNR							x	
<i>Atalopedes campestris</i>	Sachem	SNA					x			
<i>Atteva aurea</i>	Ailanthus Webworm Moth	SNR						x		
<i>Battus philenor</i>	Pipevine Swallowtail	SNA					x			
<i>Boloria bellona</i>	Meadow Fritillary	S5					x			
<i>Catocala cerogama</i>	Yellow-banded Underwing	S5						x		
<i>Catocala minuta</i>	Little Underwing	SNR							x	
<i>Catocala parta</i>	Mother Underwing	S5							x	
<i>Catocala piatrix</i>	The Penitent	S4							x	
<i>Celastrina lucia</i>	Northern Spring Azure	S5					x			
<i>Celastrina neglecta</i>	Summer Azure	S5					x			
<i>Ceratomia undulosa</i>	Waved Sphinx Moth	S5							x	
<i>Cercyonis pegala</i>	Common Wood-Nymph	S5					x			
<i>Chlosyne nycteis</i>	Silvery Checkerspot	S5					x			
<i>Cisseps fulvicollis</i>	Yellow-collared Scape Moth	SNR							x	
<i>Coenonympha tullia</i>	Common Ringlet	S5					x			

Species Name		Conservation Rank				Source				
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	NHIC ⁽¹⁾	Ontario Butterfly Atlas ⁽²⁾	iNaturalist ⁽³⁾	Ontario Moth Atlas ⁽⁴⁾	Matrix Field Observations
<i>Colias eurytheme</i>	Orange Sulphur	S5					x			
<i>Colias philodice</i>	Clouded Sulphur	S5					x	x		
<i>Ctenucha virginica</i>	Virginia Ctenucha Moth	S5							x	
<i>Cupido comyntas</i>	Eastern Tailed Blue	S5					x	x		
<i>Danaus plexippus</i>	Monarch	S2N,S4B	SC	END	SC		x	x		
<i>Darapsa myron</i>	Hog Sphinx	SU							x	
<i>Datana integerrima</i>	Walnut Caterpillar Moth	SNR							x	
<i>Deidamia inscripta</i>	Lettered Sphinx	SNA							x	
<i>Epargyreus clarus</i>	Silver-spotted Skipper	S4					x			
<i>Erynnis baptisiae</i>	Wild Indigo Duskywing	S4					x			
<i>Erynnis juvenalis</i>	Juvenal's Duskywing	S5					x			
<i>Estigmene acrea</i>	Salt Marsh Moth	S5							x	
<i>Euchaetes egle</i>	Milkweed Tussock Moth	S4?							x	
<i>Eumorpha pandorus</i>	Pandorus Sphinx	S4							x	
<i>Euphyes dion</i>	Dion Skipper	S4					x			
<i>Euphyes vestris</i>	Dun Skipper	S5					x			
<i>Euptoieta claudia</i>	Variegated Fritillary	SNA					x			
<i>Feniseca tarquinius</i>	Harvester	S4					x			
<i>Glaucopsyche lygdamus</i>	Silvery Blue	S5					x			
<i>Halysidota tessellaris</i>	Banded Tussock Moth	S5						x	x	
<i>Haploa confusa</i>	Confused Haploa	S5							x	
<i>Hemaris diffinis</i>	Snowberry Clearwing Moth	S4S5							x	
<i>Hyalophora cecropia</i>	Cecropia Moth	S5						x	x	
<i>Hylephila phyleus</i>	Fiery Skipper	SNA					x			
<i>Junonia coenia</i>	Common Buckeye	SNA					x			
<i>Leptotes marina</i>	Marine Blue	SNA					x			
<i>Lethe anhedon</i>	Northern Pearly-Eye	S5					x			
<i>Lethe eurydice</i>	Eyed Brown	S5					x			
<i>Libytheana carinenta</i>	American Snout	SNA					x			
<i>Limenitis archippus</i>	Viceroy	S5					x			
<i>Limenitis arthemis arthemis</i>	White Admiral	S5					x			
<i>Limenitis arthemis astyanax</i>	Red-spotted Purple	S5					x			
<i>Lophocampa caryae</i>	Hickory Tussock Moth	SNR						x	x	
<i>Lophocampa maculata</i>	Spotted Tussock Moth	S4							x	
<i>Lymantria dispar</i>	Gypsy Moth	SNA						x		
<i>Malacosoma americana</i>	Eastern Tent Caterpillar Moth	S5						x		
<i>Megisto cymela</i>	Little Wood-Satyr	S5					x			
<i>Noctua pronuba</i>	Large Yellow Underwing Moth	SNA						x		
<i>Nymphalis antiopa</i>	Mourning Cloak	S5					x	x		
<i>Nymphalis l-album</i>	Compton Tortoiseshell	S5					x			
<i>Paonias excaecata</i>	Blinded Sphinx	S5							x	
<i>Papilio cressphontes</i>	Giant Swallowtail	S4					x			
<i>Papilio glaucus</i>	Eastern Tiger Swallowtail	S5					x	x		

Species Name		Conservation Rank				Source				
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	NHIC ⁽¹⁾	Ontario Butterfly Atlas ⁽²⁾	iNaturalist ⁽³⁾	Ontario Moth Atlas ⁽⁴⁾	Matrix Field Observations
<i>Papilio polyxenes</i>	Black Swallowtail	S5					x	x		
<i>Peridroma saucia</i>	Variegated Cutworm Moth	S5						x		
<i>Phigalia titea</i>	The Half-wing	SNR						x		
<i>Phoebis sennae</i>	Cloudless Sulphur	SNA					x			
<i>Pholisora catullus</i>	Common Sootywing	S4					x			
<i>Phragmatobia fuliginosa</i>	Ruby Tiger Moth	S4?							x	
<i>Phyciodes cocyta</i>	Northern Crescent	S5					x			
<i>Phyciodes tharos</i>	Pearl Crescent	S4					x	x		
<i>Pieris oleracea</i>	Mustard White	S4					x			
<i>Pieris rapae</i>	Cabbage White	SNA					x	x		
<i>Pieris virginiensis</i>	West Virginia White	S3	SC				x			
<i>Poanes hobomok</i>	Hobomok Skipper	S5					x			
<i>Poanes viator</i>	Broad-winged Skipper	S4					x			
<i>Polites mystic</i>	Long Dash Skipper	S5					x			
<i>Polites origenes</i>	Crossline Skipper	S4					x			
<i>Polites peckius</i>	Peck's Skipper	S5					x	x		
<i>Polites themistocles</i>	Tawny-edged Skipper	S5					x			
<i>Polites vibex</i>	Whirlabout	SNA					x			
<i>Polygonia comma</i>	Eastern Comma	S5					x			
<i>Polygonia interrogationis</i>	Question Mark	S5					x	x		
<i>Polyommatus icarus</i>	European Common Blue	SNA					x			
<i>Pompeius verna</i>	Little Glassywing	S4					x			
<i>Pyralis farinalis</i>	Meal Moth	SNR						x		
<i>Pyrisitia lisa</i>	Little Yellow	SNA					x			
<i>Pyrrharctia isabella</i>	Isabella Tiger Moth	S5						x	x	
<i>Satyrium acadica</i>	Acadian Hairstreak	S4					x			
<i>Satyrium calanus</i>	Banded Hairstreak	S4					x			
<i>Satyrium caryaevorus</i>	Hickory Hairstreak	S4					x			
<i>Satyrium edwardsii</i>	Edwards' Hairstreak	S4					x			
<i>Satyrium liparops</i>	Striped Hairstreak	S5					x			
<i>Speyeria cybele</i>	Great Spangled Fritillary	S5					x			
<i>Sphexcodina abbottii</i>	Abbott's Sphinx	S4							x	
<i>Spilosoma virginica</i>	Virginian Tiger Moth	S5							x	
<i>Thorybes pylades</i>	Northern Cloudywing	S5					x			
<i>Thymelicus lineola</i>	European Skipper	SNA					x			
<i>Tolype notialis</i>	Small Tolyte	SNR							x	
<i>Tolype velleda</i>	Large Tolyte	SNR							x	
<i>Vanessa atalanta</i>	Red Admiral	S5					x	x		
<i>Vanessa cardui</i>	Painted Lady	S5					x			
<i>Vanessa virginiensis</i>	American Lady	S5					x			
<i>Wallengrenia egeremet</i>	Northern Broken-Dash	S5					x			
<i>Zerene cesonia</i>	Southern Dogface	SNA					x			
Mantodea	Mantises									

Species Name		Conservation Rank				Source				
Scientific Name	Common Name	Provincial (S-rank)	Provincial (ESA)	National (COSEWIC)	National (SARA)	NHIC ⁽¹⁾	Ontario Butterfly Atlas ⁽²⁾	iNaturalist ⁽³⁾	Ontario Moth Atlas ⁽⁴⁾	Matrix Field Observations
<i>Mantis religiosa</i>	Praying Mantis	SNA						x		
<i>Odonata</i>	Damselflies and Dragonflies									
<i>Calopteryx maculata</i>	Ebony Jewelwing	S5						x		x
<i>Ischnura verticalis</i>	Eastern Forktail	S5						x		
<i>Orthoptera</i>	Grasshoppers, Katydid, Crickets, and related insects									
<i>Dissosteira carolina</i>	Carolina Grasshopper	S4S5						x		
<i>Gryllus pennsylvanicus</i>	Fall Field Cricket	S5						x		
<i>Meconema thalassinum</i>	Drumming Katydid	SNA						x		
<i>Microcentrum rhombifolium</i>	Greater Angle-wing Katydid	S4						x		
Total						0	70	48	28	1

Notes:

S-rank

- S1 - Critically Imperiled
- S2 - Imperiled
- S3 - Vulnerable
- S4 - Apparently Secure
- S5 - Secure
- SU - Unrankable
- SNA - Unranked
- SX - Presumed Extirpated
- SH - Possibly Extirpated
- S#? - Rank Uncertain

COSEWIC

- NAR - Not at Risk
- SC - Special Concern
- THR - Threatened
- END - Endangered
- EXT - Extinct
- EXP - Extirpated
- DD - Data Deficient

SARA Schedule

- Schedule 1 - Officially protected under SARA
- Schedule 2 - threatened/endangered; may be reassessed for consideration for inclusion to Schedule 1
- Schedule 3 - special concern; may be reassessed for consideration for inclusion to Schedule 1

ESA

- SC - Special Concern
- THR - Threatened
- END - Endangered
- EXT - Extinct
- EXP - Extirpated

COSSARO

- NAR - Not at Risk
- SC - Special Concern
- THR - Threatened
- END - Endangered
- EXP - Extirpated
- DD - Data Deficient

Additional Notes

- ESA - *Endangered Species Act*
- COSEWIC - Committee on the Status of Endangered Wildlife in Canada
- SARA - *Species at Risk Act*
- NHIC - Natural Heritage Information Centre

Sources:

- 1 Ontario Ministry of Natural Resources and Forestry (MNRF). 2024. Make a Map: Natural Heritage Areas. Mapping application. Accessed March 2024. https://www.lioapplications.lrc.gov.on.ca/Natural_Heritage/index.html?viewer=Natural_Heritage.Natural_Heritage&locale=en-CA
- 2 Toronto Entomologists' Association (TEA). 2024. Ontario Butterfly Atlas. Accessed March 2024. <https://www.ontarioinsects.org/atlas/>
- 3 iNaturalist (iNaturalist). 2024. Observations. Accessed March 2024. <https://www.inaturalist.org/observations>
- 4 Toronto Entomologists' Association (TEA). 2024. Ontario Butterfly Atlas. Accessed March 2024. <https://www.ontarioinsects.org/moth/index.html>

TABLE 5 Mollusc Species

Species		Conservation Rank				Source			
Scientific Name	Common Name	Provincial (S-RANK)	Provincial (ESA)	National (COSEWIC)	National (SARA)	NHIC (1)	iNaturalist (2)	DFO (3)	Matrix Field Observations
Unionidae									
<i>Amblema plicata</i>	Three-ridge	S4					x		
Arionidae									
<i>Arion subfuscus</i>	Dusky Arion Slug	SNA					x		
Helicidae									
<i>Cepaea nemoralis</i>	Grovesnail	SNA					x		
TOTAL:						0	3	0	0

Sources:

- 1 Ontario Ministry of Natural Resources and Forestry (MNRF). 2024. Make a Map: Natural Heritage Areas. Mapping application. Accessed March 2024. https://www.lioapplications.lrc.gov.on.ca/Natural_Heritage/index.html?viewer=Natural_Heritage.Na
- 2 iNaturalist. 2024. Observations. Accessed March 2024. <https://www.inaturalist.org/observations>
- 3 Fisheries and Oceans Canada (DFO). 2024. Aquatic Species at Risk Map. Accessed March 2024. <https://www.dfo-mpo.gc.ca/species-especes/sara-lep/map-carte/index-eng.html>

Table 6 Mammal Species

Species Name		Conservation Ranking				Source				
Scientific Name	Common Name	S-RANK	ESA	COSEWIC	SARA	NHIC (1)	Ontario Mammals (2)	iNaturalist (3)	MECP Information Request (2021)	Matrix Field Observations
Artiodactyla		Deer and Bison								
<i>Odocoileus virginianus</i>	White-tailed Deer	S5					x			
Carnivora		Carnivores								
<i>Canis latrans</i>	Coyote	S5					x	x		x
<i>Lontra canadensis</i>	North American River Otter	S5					x			
<i>Lynx rufus</i>	Bobcat	S4					x			
<i>Martes americana</i>	American Marten	S5					x			
<i>Mephitis mephitis</i>	Striped Skunk	S5					x	x		
<i>Mustela frenata</i>	Long-tailed Weasel	SU					x			
<i>Mustela nivalis</i>	Least Weasel	SU					x			
<i>Neogale vison</i>	American Mink	S4					x			
<i>Procyon lotor</i>	Northern Raccoon	S5					x	x		x
<i>Ursus americanus</i>	American Black Bear	S5					x			
<i>Vulpes vulpes</i>	Red Fox	S5					x	x		
Chiroptera		Bats								
<i>Eptesicus fuscus</i>	Big Brown Bat	S4					x	x		
<i>Lasionycteris noctivagans</i>	Silver-haired Bat	S4					x			
<i>Lasiurus borealis</i>	Red Bat	S4					x			
<i>Lasiurus cinereus</i>	Hoary Bat	S4					x			
<i>Myotis leibii</i>	Eastern Small-footed Myotis	S2S3	END				x		x	
<i>Myotis lucifugus</i>	Little Brown Myotis	S3	END	END	END		x		x	
<i>Myotis septentrionalis</i>	Northern Myotis	S3	END	END	END		x		x	
<i>Perimyotis subflavus</i>	Tricolored Bat	S3?	END	END	END		x			
Didelphimorphia		Opposums								
<i>Didelphis virginiana</i>	Virginia Opossum	S4					x	x		
Lagomorpha		Rabbits and Hares								
<i>Sylvilagus floridanus</i>	Eastern Cottontail	S5					x	x		
<i>Lepus americanus</i>	Snowshoe Hare	S5					x			
Rodentia		Rodents								
<i>Castor canadensis</i>	Beaver	S5					x	x		
<i>Erethizon dorsatum</i>	Porcupine	S5					x			
<i>Glaucomys sabrinus</i>	Northern Flying Squirrel	S5					x			
<i>Glaucomys volans</i>	Southern Flying Squirrel	S4					x			
<i>Marmota monax</i>	Woodchuck	S5					x			
<i>Microtus pennsylvanicus</i>	Meadow Vole	S5					x			
<i>Mus musculus</i>	House Mouse	SNA					x			
<i>Myodes gapperi</i>	Southern Red-backed Vole	S5					x			
<i>Napaeozapus insignis</i>	Woodland Jumping Mouse	S5					x			
<i>Ondatra zibethicus</i>	Muskrat	S5					x			

Scientific Name	Common Name	S-RANK	ESA	COSEWIC	SARA	NHIC (1)	Ontario Mammals (2)	iNaturalist (3)	MECP Information Request (2021)	Matrix Field Observations
<i>Peromyscus leucopus</i>	White-footed Mouse	S5					x			
<i>Peromyscus maniculatus</i>	Deer Mouse	S5					x			
<i>Rattus norvegicus</i>	Norway Rat	SNA					x			
<i>Sciurus carolinensis</i>	Grey Squirrel	S5					x	x		x
<i>Synaptomys cooperi</i>	Southern Bog Lemming	S4					x			
<i>Tamias striatus</i>	Eastern Chipmunk	S5					x			
<i>Tamiasciurus hudsonicus</i>	Red Squirrel	S5					x			
<i>Zapus hudsonius</i>	Meadow Jumping Mouse	S5					x			
Soricomorpha										
<i>Blarina brevicauda</i>	Northern Short-tailed Shrew	S5					x			
<i>Condylura cristata</i>	Star-nosed Mole	S5					x			
<i>Parascalops breweri</i>	Hairy-tailed Mole	S4					x			
<i>Sorex cinereus</i>	Masked Shrew	S5					x			
<i>Sorex hoyi</i>	Pygmy Shrew	S4					x			
TOTAL:						0	46	9	3	3

Sources:

1 Ontario Ministry of Natural Resources and Forestry (MNRF). 2024. Make a Map: Natural Heritage Areas. Mapping application. Accessed March 2024. https://www.lioapplications.lrc.gov.on.ca/Natural_Heritage/index.html?viewer=Natura

2 iNaturalist. 2024. Ontario Mammals. Accessed March 2024. <https://www.inaturalist.org/guides/1327?view=card>

3 iNaturalist. 2024. Observations. Accessed March 2024. <https://www.inaturalist.org/observations>

COMMON NAME	BOTANICAL NAME	COEFFICIENT OF CONSERVATISM	WETNESS INDEX	WEEDINESS INDEX	INVASIVE SPECIES ONTARIO	PROVINCIAL RANK	ESA STATUS	COSEWIC STATUS (2016-08-19)	SARA STATUS (2016-08-19)	GLOBAL RANK	REGIONAL STATUS 7E - CAROLINIAN ZONE - 2017	LOCAL STATUS CVC and PEEL	All Species	TRCA Data Request	iNaturalist	MECP Information Request (2021)	NHIC	STUDY AREA MATRIX FIELD OBSERVATIONS
	Reference											Credit Valley Conservation, 2002						
FERNS & ALLIES	PTERIDOPHYTES												x	o	o	o		x
Wood Fern Family	Dryopteridaceae												x	o	o			x
Ostrich Fern	<i>Matteuccia struthiopteris</i>	5	0			S5				G5	C		x		x			
Horsetail Family	Equisetaceae												x	o	o			x
Field Horsetail	<i>Equisetum arvense</i>	0	0			S5				G5	C		x		x			
CONIFERS	GYMNOSPERMS												x	o	o			
Cedar Family	Cupressaceae												x	o	o			
Eastern White Cedar	<i>Thuja occidentalis</i>	4	-3			S5				G5	C		x	x				x
Pine Family	Pinaceae												x	o	o			x
Balsam Fir	<i>Abies balsamea</i>	5	-3			S5				G5	R		x					x
White Spruce	<i>Picea glauca</i>	6	3			S5				G5	U		x	x	x			x
Red Pine	<i>Pinus resinosa</i>	8	3			S5				G5	R	R	x	x				x
Eastern White Pine	<i>Pinus strobus</i>	4	3			S5				G5	C		x	x				x
Eastern Hemlock	<i>Tsuga canadensis</i>	7	3			S5				G5	C		x	x	x			x
DICOTS	DICOTYLEDONS												x	o	o	o		
Maple Family	Aceraceae												x		o			x
Hedge Maple	<i>Acer campestre</i>		5			SNA				GNR	IR		x		x			
Amur Maple	<i>Acer ginnala</i>		5	-2	4	SNA				G-TNR			x					x
Manitoba Maple	<i>Acer negundo</i>	0	0		1	S5				G5	C		x		x			x
Norway Maple	<i>Acer platanoides</i>		5	-3	2	SNA				GNR	IU		x					x
Silver Maple	<i>Acer saccharinum</i>	5	-3			S5				G5	C		x					x
Tatarian Maple	<i>Acer tataricum</i>										IR		x					x
Sumac or Cashew Family	Anacardiaceae												x		o			x
Staghorn Sumac	<i>Rhus typhina</i>	1	3			S5				G5	C		x		x			x
Carrot or Parsley Family	Apiaceae												x	o	o			x
Canadian Honewort	<i>Cryptotaenia canadensis</i>	5	0			S5				G5	C		x		x			
Wild Carrot	<i>Daucus carota</i>		5	-2		SNA				GNR	IC		x		x			x
Wild Parsnip	<i>Pastinaca sativa</i>		5	-3		SNA				GNR	IU		x		x			x
Dogbane Family	Apocynaceae												x		o			x
Common Periwinkle	<i>Vinca minor</i>		5	-2	2	SNA				GNR	IX		x		x			x
Ginseng Family	Araliaceae												x		o			x
English Ivy	<i>Hedera helix</i>					SNA					IR		x		x			
Milkweed Family	Asclepiadaceae												x		o			x
Common Milkweed	<i>Asclepias syriaca</i>	0	5			S5				G5	C		x		x			x
European Swallow-wort	<i>Vincetoxicum rossicum</i>		5	-3	1	SNA				GNR	IX		x		x			x
Composite or Aster Family	Asteraceae												x	o	o	o		x
Common Yarrow	<i>Achillea millefolium</i>		3	-1		SNA				G5	IX		x		x			
Common Ragweed	<i>Ambrosia artemisiifolia</i>	0	3			S5				G5	C		x		x			
Giant Ragweed	<i>Ambrosia trifida</i>	0	0			S5				G5	C		x		x			
Great Burdock	<i>Arctium lappa</i>		3			SNA				GNR	IU		x		x			
Common Burdock	<i>Arctium minus</i>		3	-2		SNA				GNR	IC		x					x
Common Wormwood	<i>Artemisia absinthium</i>		5	-1	3	SNA				GNR	IR		x		x			
Heart-leaved Aster	<i>Symphotrichum cordifolium</i>	5	5			S5				G5	C		x		x			
Panicled Aster	<i>Symphotrichum lanceolatum</i>	3	-3			S5				G5	C		x					x
Calico Aster	<i>Symphotrichum lateriflorum</i>	3	0			S5				G5	C		x					x
New England Aster	<i>Symphotrichum novae-angliae</i>	2	-3			S5				G5	C		x					x
Chicory	<i>Cichorium intybus</i>		5	-1		SNA				GNR	IC		x		x			x

COMMON NAME	BOTANICAL NAME	COEFFICIENT OF CONSERVATISM	WETNESS INDEX	WEEDINESS INDEX	INVASIVE SPECIES ONTARIO	PROVINCIAL RANK	ESA STATUS	COSEWIC STATUS (2016-08-19)	SARA STATUS (2016-08-19)	GLOBAL RANK	REGIONAL STATUS 7E - CAROLINIAN ZONE - 2017	LOCAL STATUS CVC and PEEL	All Species	TRCA Data Request	iNaturalist	MECP Information Request (2021)	NHIC	STUDY AREA MATRIX FIELD OBSERVATIONS
Canada Thistle	<i>Cirsium arvense</i>		3	-1	1	SNA				G5	IC		x		x			x
Bull Thistle	<i>Cirsium vulgare</i>		3	-1		SNA				GNR	IC		x		x			
Philadelphia Fleabane	<i>Erigeron philadelphicus</i>	1	-3			S5				G5	C		x		x			x
Tall Boneset	<i>Eupatorium altissimum</i>	3	5			S4				G5	R		x		x			
White Snakeroot	<i>Ageratina altissima</i>					S5				G5	C		x		x			
Common Sunflower	<i>Helianthus annuus</i>		3	-1		SNA				G5	IR		x		x			
Nipplewort	<i>Lapsana communis</i>		3	-2	4	SNA				GNR	IX		x		x			x
Black-eyed Susan	<i>Rudbeckia hirta</i>	0	3			S5				G5	C		x					x
Brown-eyed Susan	<i>Rudbeckia triloba</i>		3	-1		SNA				G5	IX		x		x			
Tall Goldenrod	<i>Solidago altissima</i>	1	3			S5				G5	C		x					x
Canada Goldenrod	<i>Solidago canadensis var. canadensis</i>	1	3			S5				G5	C		x					x
Canada Goldenrod	<i>Solidago canadensis var. hargeri</i>										U		x					x
Zig-zag Goldenrod	<i>Solidago flexicaulis</i>	6	3			S5				G5	C		x		x			x
Canada Goldenrod	<i>Solidago lepida</i>	1	3			S47				G5			x					x
Common Sow-thistle	<i>Sonchus oleraceus</i>		3	-1		SNA				GNR	IX		x		x			
Common Tansy	<i>Tanacetum vulgare</i>		5	-1		SNA				GNR	IX		x		x			x
Common Dandelion	<i>Taraxacum officinale</i>		3	-2		SNA				G5	IC		x		x			x
Jack go to bed at noon	<i>Tragopogon pratensis</i>		5	-1		SNA				GNR	IX		x		x			
Coltsfoot	<i>Tussilago farfara</i>		3	-2		SNA				GNR	IC		x		x			
Rough Cocklebur	<i>Xanthium strumarium</i>	2	0			S5				G5	C		x		x			
Touch-me-not Family	Balsaminaceae												x		o			x
Jewelweed	<i>Impatiens capensis</i>	4	-3			S5				G5	C		x					x
Pale Touch-me-not	<i>Impatiens pallida</i>	7	-3			S4				G5	C	R	x		x			
Barberry Family	Berberidaceae												x	o	o			x
May-apple	<i>Podophyllum peltatum</i>	5	3			S5				G5	C		x		x			
Birch Family	Betulaceae												x	o	o			x
Paper Birch	<i>Betula papyrifera</i>	2	2			S5				G5	C		x	x				x
Blue Beech	<i>Carpinus caroliniana</i>										C		x					x
Blue Beech	<i>Carpinus caroliniana ssp. virginiana</i>	6	0			S5				G5T5			x					x
Bigonia Family	Bigoniaceae												x		o			x
Northern Catalpa	<i>Catalpa speciosa</i>		3	-1		SNA				G47	IR		x		x			
Borage Family	Boraginaceae												x		o			x
Hound's-tongue	<i>Cynoglossum officinale</i>		5	-1		SNA				GNR	IU		x		x			
Viper's Bugloss	<i>Echium vulgare</i>		5	-2		SNA				GNR	IC		x		x			
Virginia Stickweed	<i>Hackelia virginiana</i>	5	1			S5				G5	C		x					x
Field Forget-me-not	<i>Myosotis arvensis</i>		0	-1		SNA				GNR	IR		x					x
True Forget-me-not	<i>Myosotis scorpioides</i>		-5	-1	4	SNA				G5	IX		x		x			
Strict Forget-me-not	<i>Myosotis stricta</i>		5	-1		SNA				GNR	IX		x		x			
Woodland Forget-me-not	<i>Myosotis sylvatica</i>		5	-1		SNA				G5	IR		x		x			
Lungwort	<i>Pulmonaria officinalis</i>					SNA				GNR	IR		x		x			
Mustard Family	Brassicaceae												x	o	o	o		x
Garlic Mustard	<i>Alliaria petiolata</i>		0	-3	1	SNA				GNR	IC		x		x			x
Garden Yellowrocket	<i>Barbarea vulgaris</i>		0	-1	3	SNA				GNR	IC		x					x
Cut-leaved Toothwort	<i>Cardamine concatenata</i>	6	3			S5				G5	C		x		x			
Narrowleaf Bitter-cress	<i>Cardamine impatiens</i>		5	-1		SNA				GNR	IR		x		x			x
Dame's Rocket	<i>Hesperis matronalis</i>		5	-3	1	SNA				G4G5	IC		x		x			x
Annual Honesty	<i>Lunaria annua</i>		5	-1		SNA				GNR	IR		x		x			
Field Penny-cress	<i>Thlaspi arvense</i>		5	-1		SNA				GNR	IC		x		x			
Bellflower Family	Lobelia												x		o			x
Creeping Bellflower	<i>Campanula rapunculoides</i>		5	-2	4	SNA				GNR	IU		x		x			x
Honeysuckle Family	Caprifoliaceae												x	o	o	o		x
Tartarian Honeysuckle	<i>Lonicera tatarica</i>		3	-3	1	SNA				GNR	IC		x					x
Western Snowberry	<i>Symphoricarpos occidentalis</i>		5	-1		SNA				G5	IR		x					x
European Cranberrybush	<i>Viburnum opulus</i>		0	-1		SNA				G5			x		x			
Pink Family	Caryophyllaceae												x	o	o	o		x
Bladder Campion	<i>Silene latifolia</i>		5	-2		SNA				GNR	IX		x		x			
Goosefoot Family	Chenopodiaceae												x	o	o	o		x
Lamb's Quarters	<i>Chenopodium album</i>					SNA				G5	IC		x		x			
Morning-glory Family	Convolvulaceae												x		o			x
Field Bindweed	<i>Convolvulus arvensis</i>		5	-1	3	SNA				GNR	IC		x					x
Dogwood Family	Cornaceae												x		o			x
Alternate-leaved Dogwood	<i>Cornus alternifolia</i>	6	5			S5				G5	C		x					x

COMMON NAME	BOTANICAL NAME	COEFFICIENT OF CONSERVATISM	WETNESS INDEX	WEEDINESS INDEX	INVASIVE SPECIES ONTARIO	PROVINCIAL RANK	ESA STATUS	COSEWIC STATUS (2016-08-19)	SARA STATUS (2016-08-19)	GLOBAL RANK	REGIONAL STATUS 7E - CAROLINIAN ZONE - 2017	LOCAL STATUS CVC and PEEL	All Species	TRCA Data Request	iNaturalist	MECP Information Request (2021)	NHIC	STUDY AREA MATRIX FIELD OBSERVATIONS
Gray Dogwood	<i>Cornus racemosa</i>	2	-2			S5				G5	C		x					x
Red-osier Dogwood	<i>Cornus sericea</i>	2	-3			S5				G5	C				x			
Stonecrop Family	Crassulaceae												x		0			x
Goldmoss Stonecrop	<i>Sedum acre</i>		5	-3	2	SNA				GNR	IX		x		x			
Gourd Family	Cucurbitaceae												x		0			
Wild Cucumber	<i>Echinocystis lobata</i>	3	-2			S5				G5	C		x		x			
Teasel Family	Dipsacaceae												x		0			
Fuller's Teasel	<i>Dipsacus fullanum</i>		5	-1	3	SNA				GNR	IC		x		x			
Spurge Family	Euphorbiaceae												x	0	0	0		x
Three-seeded Mercury	<i>Acalypha rhomboides</i>	0	3			S5				G5	C		x		x			
Spotted Spurge	<i>Euphorbia maculata</i>		4	-1		SNA				G5?	IC		x		x			
Pea Family	Fabaceae												x	0	0	0		x
Crown-vetch	<i>Securigera varia</i>		5	-2	1	SNA				GNR	IX		x		x			
Honey Locust	<i>Gleditsia triacanthos</i>	3	0			S2?				G5	R		x					x
Bird's-foot Trefoil	<i>Lotus corniculatus</i>		1	-2	2	SNA				GNR	IC		x		x			x
Black Medick	<i>Medicago lupulina</i>		1	-1	4	SNA				GNR	IC		x					x
White Sweet-clover	<i>Mellilotus albus</i>		3	-3	2	SNA				G5	IC		x					x
Black Locust	<i>Robinia pseudoacacia</i>		4	-3	2	SNA				G5	IC		x		x			
Red Clover	<i>Trifolium pratense</i>		2	-2	4	SNA				GNR	IC		x		x			
White Clover	<i>Trifolium repens</i>		2	-1	4	SNA				GNR	IC		x		x			
Cow Vetch	<i>Vicia cracca</i>		5	-1	2	SNA				GNR	IX		x		x			x
Beech Family	Fagaceae												x	0	0	0		x
Bur Oak	<i>Quercus macrocarpa</i>	5	1			S5				G5	C		x		x			x
Red Oak	<i>Quercus rubra</i>	6	3			S5				G5	C		x	x				x
Geranium Family	Geraniaceae												x	0	0	0		x
Spotted Geranium	<i>Geranium maculatum</i>	6	3			S5				G5	C		x		x			
Herb-robert	<i>Geranium robertianum</i>		5	-2		S5				G5	C		x		x			x
Water-leaf Family	Hydrophyllaceae												x	0	0	0		x
Virginia Water-leaf	<i>Hydrophyllum virginianum</i>	6	-2			S5				G5	C		x		x			
Walnut Family	Juglandaceae												x	0	0	0		x
Butternut	<i>Juglans cinerea</i>	6	2			S2?	END	END	END	G4	U		x	x	x	x		x
Black Walnut	<i>Juglans nigra</i>	5	3			S4?				G5	C		x		x			x
Mint Family	Lamiaceae												x	0	0	0		x
Common Bugle	<i>Ajuga reptans</i>		5	-1	4	SNA				GNR	IR		x		x			
Ground Ivy	<i>Glechoma hederacea</i>		5	-2	4	SNA				GNR	IC		x		x			x
Purple Dead-nettle	<i>Lamium purpureum</i>		5	-2		SNA				GNR	IR		x		x			
Common Motherwort	<i>Leonurus cardiaca</i>												x		x			x
Common Motherwort	<i>Leonurus cardiaca ssp. cardiaca</i>		5	-2		SNA				GNRTNR			x					x
Catnip	<i>Nepeta cataria</i>		1	-2	4	SNA				GNR	IC		x		x			x
Loosestrife Family	Lythraceae												x	0	0	0		x
Purple Loosestrife	<i>Lythrum salicaria</i>		-5	-3	1	SNA				G5	IC		x		x			
MoONSEED Family	Menispermaceae												x	0	0	0		
Common Moonseed	<i>Menispermum canadense</i>	7	0			S4				G5	C	R	x	x				
Mulberry Family	Moraceae												x		0			x
White Mulberry	<i>Morus alba</i>		0	-3	1	SNA				GNR	IC		x		x			x
Olive Family	Oleaceae												x		0			x
White Ash	<i>Fraxinus americana</i>	4	3			S4				G5	C		x					x
Green Ash	<i>Fraxinus pennsylvanica</i>	3	-3			S4				G5	C		x					x
Common Lilac	<i>Syringa vulgaris</i>		5	-2	2	SNA				GNR	IX		x		x			x
Wood Sorrel Family	Oxalidaceae												x		0			x
Creeping Wood-sorrel	<i>Oxalis corniculata</i>		3	-1		SNA				GNR	IR		x		x			
Common Yellow Oxalis	<i>Oxalis stricta</i>	0	3			S5				G5	C		x					x
Poppy Family	Papaveraceae												x		0			x
Celandine	<i>Chelidonium majus</i>		5	-3		SNA				GNR	IU		x		x			
Bloodroot	<i>Sanguinaria canadensis</i>	5	4			S5				G5	C		x		x			x
Plantain Family	Plantaginaceae												x		0			x
Common Plantain	<i>Plantago major</i>		-1	-1		SNA				G5	IC		x		x			x
Smartweed Family	Polygonaceae												x	0	0			x
Hedge Bindweed	<i>Fallopia dumetorum</i>												x		x			x
Japanese Knotweed	<i>Fallopia japonica</i>		3	-1	2	SNA				G?	IX		x					x
Japanese Knotweed	<i>Fallopia japonica var. japonica</i>												x					x
Primrose Family	Primulaceae												x	0	0			x
Fringed Loosestrife	<i>Lysimachia ciliata</i>	4	-3			S5				G5	C		x		x			x
Creeping Jenny	<i>Lysimachia nummularia</i>		-4	-3	2	SNA				GNR	IC		x					x
Buttercup Family	Ranunculaceae												x	0	0			x
Tall Buttercup	<i>Ranunculus acris</i>		-2	-2		SNA				G5	IC		x					x
Buckthorn Family	Rhamnaceae												x	0	0			x
Common Buckthorn	<i>Rhamnus cathartica</i>		3	-3	1	SNA				GNR	IC		x		x			x
Rose Family	Rosaceae												x	0	0			x
Smooth Serviceberry	<i>Amelanchier laevis</i>	5	5			S5				G5	C		x		x			
Hawthorn species	<i>Crataegus sp.</i>	4	5										x					x
Woodland Strawberry	<i>Fragaria vesca</i>										U		x		x			x
Wild Strawberry	<i>Fragaria virginiana</i>	2	1			S5				G5	C		x					x
White Avens	<i>Geum canadense</i>	3	0			S5				G5	C		x					x
Wood Avens	<i>Geum urbanum</i>		5	-1		SNA				G5	IX		x		x			x
Siberian Crabapple	<i>Malus baccata</i>					SNA				GNR	IR		x					x
Ninebark	<i>Physocarpus opulifolius</i>	5	-2			S5				G5	U	R	x		x			

COMMON NAME	BOTANICAL NAME	COEFFICIENT OF CONSERVATISM	WETNESS INDEX	WEEDINESS INDEX	INVASIVE SPECIES ONTARIO	PROVINCIAL RANK	ESA STATUS	COSEWIC STATUS (2016-08-19)	SARA STATUS (2016-08-19)	GLOBAL RANK	REGIONAL STATUS 7E - CAROLINIAN ZONE - 2017	LOCAL STATUS CVC and PEEL	All Species	TRCA Data Request	iNaturalist	MECP Information Request (2021)	NHIC	STUDY AREA MATRIX FIELD OBSERVATIONS
Rough-fruited Cinquefoil	<i>Potentilla recta</i>		5	-2		SNA				GNR	IC		x					x
Canada Plum	<i>Prunus nigra</i>	4	4			S4				G4G5	U		x		x			
Pin Cherry	<i>Prunus pensylvanica</i>	3	4			S5				G5	C		x	x				x
Black Cherry	<i>Prunus serotina</i>	3	3			S5				G5	C		x					x
Nanking Cherry	<i>Prunus tomentosa</i>		5	-1		SNA				GNR	IR		x		x			
Choke Cherry	<i>Prunus virginiana</i>	2	1			S5				G5	C		x					x
Dog Rose	<i>Rosa canina</i>		5	-1		SNA				GNR	IX		x		x			
Multiflora Rose	<i>Rosa multiflora</i>		3	-3	1	SNA				GNR	IC		x		x			
American Red Raspberry	<i>Rubus idaeus</i>	0	-2			SNA				G5	C		x					x
Purple Flowering Raspberry	<i>Rubus odoratus</i>	3	5			S5				G5	C		x		x			
European Mountain-ash	<i>Sorbus aucuparia</i>		5	-2	4	SNA				G5	IX		x					x
Madder Family	Rubiaceae												x	0	0			x
Cleavers	<i>Galium aparine</i>	4	3			S5				G5	C	R	x					x
Willow Family	Salicaceae												x	0	0			x
Eastern Cottonwood	<i>Populus deltoides</i>										C		x					x
Eastern Cottonwood	<i>Populus deltoides ssp. deltoides</i>	4	-1			S5				G5T5	C		x					x
Trembling Aspen	<i>Populus tremuloides</i>	2	0			S5				G5	C		x		x			
Crack Willow	<i>Salix fragilis</i>		-1	-3	3	SE				GNR	IC		x					x
Saxifrage Family	Saxifragaceae												x	0	0			x
Ditch Stonecrop	<i>Penthorum sedoides</i>	4	-5			S5				G5	C		x		x			
Figwort Family	Scrophulariaceae												x	0	0			x
Butter-and-eggs	<i>Linaria vulgaris</i>		5	-1	4	SNA				GNR	IC		x		x			x
Common Mullein	<i>Verbascum thapsus</i>		5	-2		SNA				GNR	IC		x		x			
Gray Field Speedwell	<i>Veronica polita</i>		5	-1		SNA				GNR	IX		x		x			
Thyme-leaved Speedwell	<i>Veronica serpyllifolia</i>	0	-3			SNA?				G5	IC		x		x			
Nightshade Family	Solanaceae												x	0	0			x
Bittersweet Nightshade	<i>Solanum dulcamara</i>		0	-2	3	SNA				GNR	IC		x		x			x
Linden Family	Tiliaceae												x	0	0			x
American Basswood	<i>Tilia americana</i>	4	3			S5				G5	C		x					x
Little Leaf Linden	<i>Tilia cordata</i>				4	SNA				GNR	IR		x		x			
Elm Family	Ulmaceae												x	0	0			x
American Elm	<i>Ulmus americana</i>	3	-2			S5				G5	C		x					x
Siberian Elm	<i>Ulmus pumila</i>		5	-1	2	SNA				GNR	IX		x					x
Slippery Elm	<i>Ulmus rubra</i>	6	0			S5				G5	C		x	x				x
Nettle Family	Urticaceae												x		0			x
Stinging Nettle	<i>Urtica dioica ssp. dioica</i>		-1	-1	3	SNA				G5T5?	IR		x					x
Vervain Family	Verbenaceae												x		0			x
Blue Vervain	<i>Verbena hastata</i>	4	-4			S5				G5	C		x					x
Violet Family	Violaceae												x	0	0			x
Sweet Violet	<i>Viola odorata</i>		5	-1	4	SNA				GNR	IR		x		x			x
Woolly Blue Violet	<i>Viola sororia</i>	4	1			S5				G5	C		x		x			x
Grape Family	Vitaceae												x		0			x
Virginia Creeper	<i>Parthenocissus quinquefolia</i>	6	1			S4?				G5	U		x					x
Riverbank Grape	<i>Vitis riparia</i>	0	-2			S5				G5	C		x		x			x
MONOCOTS	MONOCOTYLEDONS												x	0	0			x
Arum Family	Araceae												x	0	0			x
Jack-in-the-pulpit	<i>Arisaema triphyllum</i>	5	-2			S5				G5	C		x		x			
Skunk Cabbage	<i>Symplocarpus foetidus</i>	7	-5			S5				G5	C	R	x		x			
Lily Family	Liliaceae												x	0	0			x
European Lily-of-the-valley	<i>Convallaria majalis</i>		5	-2	3	SNA				G5	IX		x		x			
Yellow Trout-lily	<i>Erythronium americanum</i>	5	5			S5				G5	C		x		x			
Snowdrop	<i>Galanthus nivalis</i>					SNA				GNR	IR		x		x			
Canada Mayflower	<i>Maianthemum canadense</i>	5	0			S5				G5	C		x	x				
Large False Solomon's Seal	<i>Maianthemum racemosum</i>	4	3			S5				G5	C		x		x			x
Starry False Solomon's Seal	<i>Maianthemum stellatum</i>	6	1			S5				G5	C		x		x			
Common Grape Hyacinth	<i>Muscari botryoides</i>		5	-1		SNA				GNR	IR		x		x			
Daffodil	<i>Narcissus pseudonarcissus</i>					SNA				GNR	IR		x					x
Sleepydicik	<i>Ornithogalum umbellatum</i>		1	-1		SNA				G3G5	IR		x		x			
Siberian Squill	<i>Scilla sibirica</i>		5	-1	2	SNA				GNR	IR		x		x			x
Grass Family	Poaceae												x	0	0			x
Smooth Brome	<i>Bromus inermis</i>		5	-3	4	SNA				G5TNR	IC		x					x
Poverty Oat Grass	<i>Danthonia spicata</i>	5	5			S5				G5	C		x	*				
Foxtail Grass	<i>Hordeum jubatum</i>					SNA				G5T5	IU		x		x			
Reed Canary Grass	<i>Phalaris arundinacea</i>	0	-4			S5				G5	C		x					x
Timothy	<i>Phleum pratense</i>		3	-1		SNA				GNR	IC		x					x
Canada Blue Grass	<i>Poa compressa</i>	0	2			SNA				GNR	IC		x					x

APPENDIX B
Site Photographs



*Matrix Supplied
May 24, 2019*

1. Most upstream extent of the study area approximately 500m upstream of Dixie Road



*Matrix Supplied
July 10, 2019*

2. Channel armoring changes from large boulders to armour stone along the banks.



*Matrix Supplied
May 24, 2019*

3. Riparian forest community along the north side of the channel upstream of Dixie Road.



*Matrix Supplied
May 24, 2019*

4. Riparian forest community along the south side of the channel upstream of Dixie Road



*Matrix Supplied
July 10, 2019*

5. The channel continues to be heavily armored moving downstream towards Dixie Road.



*Matrix Supplied
July 10, 2019*

6. There is spare amounts of fish habitat present which consists of overhanging vegetation, and some larger cobbles and boulders.



*Matrix Supplied
May 24, 2019*

7. Little Etobicoke Creek downstream of Dixie Road



*Matrix Supplied
May 24, 2019*

8. Riparian habitat north of the channel and downstream of Dixie Road.



*Matrix Supplied
July 10, 2019*

9. Channel continues to be heavily armoured downstream of Dixie Road with overhanging vegetation.



*Matrix Supplied
July 10, 2019*

10. Channel has larger areas of deposition directly downstream of Dixie Road which has created areas of concentrated flows.



*Matrix Supplied
August 22, 2019*

11. Overhanging vegetation and fallen trees over the channel



*Matrix Supplied
August 22, 2019*

12. Armourstone ends approximately 500m upstream of Dixie Road.

*Matrix Supplied
August 22, 2019*



13. Exposed banks are undercut providing cover for fish.

*Matrix Supplied
August 22, 2019*



14. Channel has larger areas of deposition directly downstream of Dixie Road which has created areas of concentrated flows.

*Matrix Supplied
August 22, 2019*



15. Butternut tree within the Study Area.



*Matrix Supplied
August 22, 2019*

16. Woody debris within the Channel.



*Matrix Supplied
August 22, 2019*

17. Channel widens at confluence with small trib.



*Matrix Supplied
August 22, 2019*

18. Channel banks are armoured at meander bend. Large boulders within channel with overhanging vegetation.

*Matrix Supplied
August 22, 2019*



19. Gabion baskets create a slow flowing pooled habitat with overhanging vegetation.



*Matrix Supplied
August 22, 2019*

20. Channel narrows moving downstream towards Dundas.

*Matrix Supplied
July 30, 2021*



21. A representative photo of a riffle downstream of Dundas St E.



*Matrix Supplied
July 30, 2021*

22. A representative photo of a pool downstream of Dundas St E.

APPENDIX C
Species at Risk and Species of Conservation
Concern Screening

TABLE 1 Species At Risk

Taxonomy	Species	ESA Status	SARA Status	COSEWIC Status	Preferred Habitat ^{1,2}	Known Species Range ^{1,2}	Source Identifying Species Record	Probability of Occurrence within Study Area	Conclusions/ Recommendations
Avian	Bank Swallow <i>Riparia riparia</i>	THR	THR Schedule 1	THR	- Requires vertical faces in sand or silt deposits; river and lake banks, active/inactive sand and gravel pits, road cuts, soil stockpiles. - Breeding sites are located close to aerial foraging areas such as grasslands, meadows, pastures, and cropland. - Large wetlands used for nocturnal roost sites during post-breeding, migration and wintering periods.	- Common across southern Ontario, especially along Lake Erie and Lake Ontario shorelines and the Saugeen River. - Sparse populations scattered across northern Ontario.	OBBA, MECP Info Request	Moderate - potential suitable vertical river banks present within study area. This species was not observed during field investigations.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Avian	Bobolink <i>Dolichonyx oryzivorus</i>	THR	THR Schedule 1	SC	- Hayfields, pastures, wet prairie, graminoid peatlands, abandoned farm fields dominated by tall grasses, no-till cropland, small-grain fields, restored surface mining sites. - Small nests are often built on the ground in dense grasses. - Typically not abundant in short-grass prairie, alfalfa, or in row crop monocultures (corn, soybean, wheat). - TPO, TPS, CUM1.	- Southern Ontario north to James Bay.	OBBA	Low - the study area lacks suitable grassland habitat for this species	Nothing further
Avian	Chimney Swift <i>Chaetura pelagica</i>	THR	THR Schedule 1	THR	- Historically included hollow trees. - More commonly found in and around urban settlements, including chimneys and other manmade structures. - Typically close to water. - TPO, CUM1, MAM, MAS, DAO, SAS1, SAM1, SAF1 adjacent to suitable nesting habitat.	- Southern Ontario north to Timmins.	NHIC, OBBA, eBird, MECP Info Request, TRCA Data Request, Matrix Field Observation	Confirmed - this species was observed flying over the study area. There are no candidate chimneys for nesting within the study area. It is anticipated that the birds use the study area for foraging.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Avian	Eastern Meadowlark <i>Sturnella magna</i>	THR	THR Schedule 1	THR	- Moderately tall grasslands; prairies, savannahs, pastures and hayfields, alfalfa fields, weedy borders of croplands, roadsides, orchards, airports, overgrown fields. - Small trees, shrubs, or fence posts used as elevated song perches. - TPO, TPS, CUM1, CUS, and MAM2.	- Southern Ontario north to Timmins, as well as Lake of the Woods area.	OBBA	Low - the study area lacks suitable grassland habitat for this species	Nothing further
Avian	Eastern Whip-poor-will <i>Antrostomus vociferans</i>	THR	THR Schedule 1	SC	- Typically a mix of open and forested areas; savannahs, open woodlands, or openings in mature deciduous, coniferous and mixed forests (commonly pine and oak forests). - Foraging habitat may include shrubby pastures or wetlands with perches. - TPS, TPW, CUW, FOD, FOC and FOM where open areas are present.	- Southern Ontario to north side of Lake Superior.	NHIC	Moderate - potential suitable forested and open habitat is present within the study area. This species was not observed during field investigations.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Avian	Golden Eagle <i>Aquila chrysaetos</i>	END	No Status	No Status	- Nest in remote, undisturbed areas, usually building their nests on ledges on a steep cliff or riverbank, but they will also use large trees if needed. - Most hunting is done near open areas such as large bogs or tundra. During migration they could be encountered anywhere, but are most frequently seen migrating west along the shores of Lake Ontario and Erie in November.	- There are only 6 extant populations within Ontario. These are located within Northern Ontario: Fort Severn, Peawanuck, Little Shagmou River, Shagmu River, Shamattawa River, Sutton Narrows, Wachi Creek, and Winsk River. - In Ontario, breeding Golden Eagles are presently known only from the Hudson Bay Lowland, although there is some evidence suggesting they once nested much further south - From Windsor northeast to Ottawa Valley/Montreal area.	eBird	Low - the study area is outside of this species' breeding range	Nothing further
Avian	Henslow's Sparrow <i>Centronyx henslowii</i>	END	END Schedule 1	END	- Open fields with tall grasses, flowering plants, and scattered shrubs; abandoned farm fields, pastures, and wet meadows. - Prefers undisturbed, extensive, dense, tall grasslands. - Avoids grazed, harvested, burned fields, or those crowded with trees and shrubs. - TPO, CUM, and MAM that are a minimum of 30 ha in size with vegetation over 30 cm in height, a thick thatch layer, and absence of woody vegetation.	- From Windsor northeast to Ottawa Valley/Montreal area.	NHIC	Low - the study area lacks suitable grassland habitat for this species	Nothing further
Avian	Least Bittern <i>Icthyophaga exilis</i>	THR	THR Schedule 1	THR	- Marshes with emergent vegetation surrounded by open water; prefers cattail marshes with a mix of open pools and channels. - Nests are built above marsh water in stands of dense vegetation. - MAS2-1, MAS3-1, SA and DAO.	- From Collingwood to Kingston as well as small pockets near Cornwall.	OBBA	Low - the study area lacks suitable marsh habitat for this species	Nothing further
Avian	Red-headed Woodpecker <i>Melanerpes erythrocephalus</i>	END	THR Schedule 1	END	- Woodlands and woodland edges, including oak and beech forests, grasslands, orchards, riparian forests, beaver ponds, burns, parks, golf courses, and cemeteries. - Dead trees used for nesting and perching. - TPS, TPW, CUW, FOD1, FOD2, FOD4-1, FOD6, FOD7, and FOD9 that are open with an abundance of dead trees.	- Woodland Caribou Provincial Park southeast to Cornwall.	NHIC	Moderate - potential suitable forested habitat present within the study area. This species was not observed during field investigations.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Flora	Butternut <i>Juglans cinerea</i>	END	END Schedule 1	END	- Deciduous forests with moist, well-drained soil of pH 5.5 to 8; commonly found along streams. - Often grows alone in sunny openings and near forest edges. - FOD and mature hedgerows.	- South of Pembroke to Port Elgin.	MECP Info Request, TRCA Info Request, Matrix Field Observation	Confirmed - this species was observed within the study area	Conduct Butternut Health Assessment to determine whether the tree is a hybrid or a pure specimen
Herpetofauna	Blanding's Turtle (Great Lakes / St. Lawrence population) <i>Emydoidea blandingii</i>	THR	END Schedule 1	END	- Shallow, nutrient-rich habitats; typically large wetlands and shallow lakes with lots of water plants. - Nesting occurs in sand, organic soil, gravel, cobblestone, and soil-filled crevices of rock outcrops. - Overwintering occurs in pools about 1 metre in depth. - SWT2, SWT3, SWD, SWM, MAS2, SAS1, SAM1, where open water is present.	- Southern Ontario north to Sudbury, with isolated reports as far north as Timmins.	ORAA	Low - the study area lacks suitable habitat for this species	Nothing further
Herpetofauna	Jefferson Salamander <i>Ambystoma jeffersonianum</i>	END	END Schedule 1	END	- Mature deciduous or mixed upland forest containing, or adjacent to, breeding ponds. - Terrestrial habitat must include small mammal burrows or rock fissures for over-wintering below the frost line. - Breeding ponds are normally ephemeral or vernal woodland pools that dry in late summer. - FOD where permanent or temporary ponds or pools are present.	- Most commonly found within the Niagara Escarpment and Carolinian forest regions.	ORAA	Low - the study area lacks ephemeral breeding ponds for this species	Nothing further
Herpetofauna	Unisexual Ambystoma (Jefferson Salamander dependent population) <i>Ambystoma laterale-(2) jeffersonianum</i>	END	END Schedule 1	END	- Mature deciduous or mixed upland forest containing, or adjacent to, breeding ponds. - Terrestrial habitat must include small mammal burrows or rock fissures for over-wintering below the frost line. - Breeding ponds are normally ephemeral or vernal woodland pools that dry in late summer.	- Southern Ontario, centred around the Niagara Escarpment.	ORAA	Low - the study area lacks ephemeral breeding ponds for this species	Nothing further
Mammals	Eastern Small-footed Myotis (Eastern Small-footed Bat) <i>Myotis leibii</i>	END	N/A	N/A	- Summer habitat includes rock outcrops, in buildings, under bridges, or in caves, mines or hollow trees. - Roosting locations are typically changed every night. - Winter hibernation occurs in caves or mines, typically drier and colder than sites selected by other bats.	- South of Georgian Bay to Lake Erie and east to the Pembroke area, the Bruce Peninsula, the Espanola area, and Lake Superior Provincial Park.	Ontario Mammals, MECP Info Request	Low - the study area lacks suitable habitat such as rock piles or fissures for this species.	Nothing further
Mammals	Little Brown Myotis (Little Brown Bat) <i>Myotis lucifugus</i>	END	END Schedule 1	END	- Large-diameter trees, attics, abandoned buildings, and barns often used for summer colonies. - Foraging occurs over water, along waterways, and forest edges, while open areas such as clearcuts or fields are typically avoided. - Hibernacula used in winter include mines and caves that are humid and remain above freezing.	- All across Ontario, concentrated in southern Ontario.	Ontario Mammals, MECP Info Request	Moderate - the study area contains snag tree habitat adjacent to a watercourse and is considered suitable habitat for this species. This species was not observed during field investigations.	Should tree removal occur, acoustic monitoring should be completed. Any tree removal should occur outside of the maternity roosting season of April 1-September 30 of any year.

Mammals	Northern Myotis (Northern Long-eared Bat) <i>Myotis septentrionalis</i>	END	END Schedule 1	END	<ul style="list-style-type: none"> - Typically within the boreal forest, under loose bark or in the cavities of trees. - Foraging occurs over water, along waterways, and forest edges, while open areas such as clearcuts or fields are typically avoided. - Overwintering occurs in cold and humid sites such as caves or mines. - FDC, FOM, FDD, SWC, SWM, and SWD where suitable roosting (i.e. cavity trees and trees with loose bark) habitat is available. 	<ul style="list-style-type: none"> - Forested areas in southern Ontario, to the north shore of Lake Superior and occasionally as far north as Moosonee, and west to Lake Nipigon. 	Ontario Mammals, MECP Info Request	<ul style="list-style-type: none"> - Moderate - the study area contains snag tree habitat adjacent to a watercourse and is considered suitable habitat for this species. This species was not observed during field investigations. 	<ul style="list-style-type: none"> - Should tree removal occur, acoustic monitoring should be completed. - Any tree removal should occur outside of the maternity roosting season of April 1-September 30 of any year.
Mammals	Tri-colored Bat <i>Perimyotis subflavus</i>	END	END Schedule 1	END	<ul style="list-style-type: none"> - Day roost and maternity colonies are formed in older forests with large-diameter trees, barns, or other structures. - Foraging occurs over water or along streams in a forest. - Winter hibernacula include caves and mines. 	<ul style="list-style-type: none"> - Southern Ontario north to Sudbury. 	Ontario Mammals	<ul style="list-style-type: none"> - Moderate - the study area contains appropriate Oak and Maple trees adjacent to a watercourse and is considered suitable habitat for this species. This species was not observed during field investigations. 	<ul style="list-style-type: none"> - Should tree removal occur, acoustic monitoring should be completed. - Any tree removal should occur outside of the maternity roosting season of April 1-September 30 of any year.

TOTAL	17
Herpetofaunas	3
Avian	9
Aquatics	0
Invertebrates	0
Flora	1
Mammals	4

ESA Status	
END	10
THR	7
TOTAL SAR	17

TABLE 2 Species of Conservation Concern

Taxonomy	Species	ESA Status	SARA Status	COSEWIC Status	Preferred Habitat ^{1,2}	Known Species Range ^{1,2}	Source Identifying Species Record	Probability of Occurrence within Study Area	Conclusions/ Recommendations
Avian	Bald Eagle <i>Haliaeetus leucocephalus</i>	SC	No Status	No Status	- Wide variety of habitats near major lakes or rivers. - Tall trees (ie, pine or poplar) typically used for nesting. - Diet consists of fish and dead animals (ie, white-tailed deer). - FOC, FOM, FOD, SWC, SWM and SWD.	- Can be found across Ontario, from US border north to Lake of the Woods.	OBBA, eBird	Low - the study area is not along a major river. No nests were observed.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Avian	Barn Swallow <i>Hirundo rustica</i>	SC	THR Schedule 1	SC	- Cup-shaped mud nests are built on human-made structures such as open barns, under bridges, and in culverts. - Preferably constructed on rough-cut wood surfaces with right angles. - Foraging habitat includes grassy fields, pastures, cropland, lake and river shorelines, cottage areas and farmyards, islands, wetlands, and tundra. - TPO, CUM1, MAM, MAS, OAO, SAS1, SAM1, and SAF1, adjacent to suitable nesting structures.	- From southern Ontario north to Hudson Bay.	NHIC, OBBA, eBird, MECP Info Request	Moderate - potential nesting structures present in the bridge and culvert within the study area. This species was not observed during field investigations.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Avian	Canada Warbler <i>Cardellina canadensis</i>	SC	THR Schedule 1	SC	- Deciduous or coniferous forests with well-developed, dense shrub layer; commonly wet or in riparian areas. - May also include stands regenerating after natural disturbances (ie, logging). - FOC3, FOC4, FOM6, FOM7, FOM8, FOD6, FOD7, FOD8, FOD9, SWC, SWM and SWD.	- All of Ontario.	eBird, MECP Info Request	Moderate - potential suitable forested habitat present within study area. This species was not observed during field investigations.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Avian	Common Nighthawk <i>Chordeiles minor</i>	SC	SC Schedule 1	SC	- Open areas with little to no ground vegetation; logged or burned areas, rock barrens, peat bogs, lakeshores, dunes, beaches, and mine tailings. - Less commonly found in cultivated fields, orchards, mine tailings, and along gravel roads and railways. - Nesting habitat is typically open and vegetation free; may include grasslands, pastures, marshes, and riverbanks. - May also include mixed and coniferous forests. - SD, BB, RB, CUM, BD, FOM, FOX and FOD with sparsely vegetated openings.	- All of Ontario except for coastal regions of James Bay and Hudson Bay.	OBBA	Low - the study area lacks suitable unvegetated habitat for this species.	Nothing further
Avian	Eastern Wood-pewee <i>Contopus virens</i>	SC	SC Schedule 1	SC	- Mid-canopy layer of forest clearings, edges of deciduous and mixed forests, early successional clearings. - FOC, FOM, FOD, SWD, SWM and CUW.	- Southern Ontario north to Sudbury.	NHIC, OBBA, eBird, MECP Info Request, Matrix Field Observation	Confirmed - this species was heard during the 2019 field studies, suitable habitat is present within the study area.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Avian	Peregrine Falcon <i>Falco peregrinus</i>	SC	No Status	No Status	- Tall, steep cliff ledges or tall buildings from 50 m to 200 m in height, close to large bodies of water. - Can be found in tundra, coastal, prairie or urban areas. - CLO.	- Breeding population centered around Lake Superior in northwest Ontario. - May be found migrating across rest of the province.	OBBA, iNaturalist, MECP Info Request	Low - the study area lacks suitable cliff habitat for this species	Nothing further
Avian	Rusty Blackbird <i>Euphagus carolinus</i>	SC	SC Schedule 1	SC	- Wet woodlands, swamps, pond edges. - Agricultural land is used for foraging. - Boreal forest is used for breeding; conifer-dominated forests adjacent to wetlands, peat bogs, sedge meadows, marshes, swamps, and beaver ponds.	- Breeding habitat spans Hudson Bay south to Orillia. - May be seen in southern Ontario during migration.	eBird	Low - the study area is outside of this species' breeding range	Nothing further
Avian	Wood Thrush <i>Hylocichla mustelina</i>	SC	THR Schedule 1	THR	- Mature deciduous and mixed forests; moist stands of trees with well-developed undergrowth. - Tall trees are used for singing perches. - Nests are built in live saplings, trees, or shrubs, especially sugar maple or American beech. - Preferably large forest mosaics. - FOD and FOM greater than 1 ha.	- Southern Ontario north to Hearst.	OBBA, eBird, MECP Info Request	Moderate - potential suitable forested habitat present within study area. This species was not observed during field investigations.	Any vegetation removal should occur outside of the breeding bird window of April 1 to August 31
Herpetofaunas	Eastern Musk Turtle (Stinkpot) <i>Sternotherus odoratus</i>	SC	SC Schedule 1	SC	- Shallow ponds, lakes, marshes and slow-moving rivers with abundant emergent vegetation and muddy bottoms. - Nesting occurs in soil, decaying vegetation, rotting wood, rock crevices, muskrat lodges, or on open ground. - Hibernation occurs in the muddy bottoms of suitable wetland habitats. - MAS, OAO, SAS, SAM, and SAF.	- South of North Bay, primarily along the eastern shoreline of Georgian Bay and north of Lake Erie and Ontario.	ORAA	Low - the study area lacks suitable habitat for this species. Record is considered historical as it is over 30 years old.	Nothing further
Herpetofaunas	Northern Map Turtle <i>Graptemys geographica</i>	SC	SC Schedule 1	SC	- Both lakes and rivers, preferably with slow-moving currents, muddy bottoms, high-quality water, and abundant vegetation. - Habitat must contain suitable basking sites such as rocks and deadheads. - Hibernation occurs at the bottom of deep, slow-moving sections of river. - OAO, SA with emergent rocks and fallen trees.	- Southern Ontario, primarily on the shores of Georgian Bay, Lake St. Clair, Lake Erie, and Lake Ontario, and along larger rivers including the Thames, Grand, and Ottawa. - Has also been recorded on Manitoulin Island and north of Timmins.	ORAA, iNaturalist	Low - the study area lacks suitable soft-bottomed aquatic habitat	Nothing further
Herpetofaunas	Snapping Turtle <i>Chelydra serpentina</i>	SC	SC Schedule 1	SC	- Shallow wetland habitats with slow-moving water and soft bottoms; ponds, sloughs, shallow bays, river edges, or slow streams. - Nesting occurs on sandy or gravel banks or man-made structures such as roads, dams, and aggregate pits. - Overwintering occurs underwater, underneath logs, sticks, or overhanging banks, deep in mud in marshy areas, or underneath floating mats of vegetation. - OAO, SA near gravelly or sandy areas.	- Primarily southern Ontario north to Timmins; also found near Thunder Bay and Kenora.	NHIC, ORAA	Low - the study area lacks suitable soft-bottomed aquatic habitat. However, it could be used as a travel corridor.	Area required for construction will be cleared of any potential turtles prior to construction and fencing will be installed to keep any turtles from entering the work zone.
Herpetofaunas	Western Chorus Frog (Great Lakes / St. Lawrence - Canadian Shield population) <i>Pseudacris triseriata</i>	No Status	THR Schedule 1	THR	- Where both terrestrial and aquatic habitats are found in close proximity with seasonally dry temporary ponds devoid of fish. - Rarely found in permanent ponds. - Terrestrial habitat primarily lowland (marshes or wooded wetlands, low shrubs and grass) with rocks, dead trees, leaves, or in loose soil or animal burrows.	- Can be found from Georgian Bay east to Eganville, and from southern Ontario to south of Algonquin Park.	ORAA	Low - the study area lacks suitable habitat for this species. Record is considered historical as it is over 30 years old.	Nothing further
Invertebrates	Monarch <i>Danaus plexippus</i>	SC	END Schedule 1	END	- Open or disturbed habitats such as roadsides, fields, wetlands, prairies, and open forests. - Trees along the north shore of the Great Lakes are used for roosting before migrating across open water. - Caterpillars are confined to meadows and open areas where milkweed grows. - AL, TP, and CUM where milkweed is present.	- South of 50° of latitude.	OBA, iNaturalist	Moderate - milkweed plants were observed on site during the field studies. This species was not observed during field investigations.	Milkweed species are recommended to be incorporated into seed mixes during any restoration efforts.
Invertebrates	West Virginia White <i>Pieris virginianis</i>	SC	N/A	N/A	- Moist, deciduous woodlots with a supply of Toothwort (the only food source of larvae).	- Southern Ontario north to Eganville as well as Manitoulin Island, eastern shore of Georgian Bay and Lake Huron. - Concentrated in western Lake Ontario region.	OBA	Low - the study area lacks suitable habitat for this species. Record is considered historical as it is over 30 years old.	Nothing further

TOTAL	15
Herpetofaunas	5
Avian	8
Aquatics	0
Invertebrates	2
Flora	0
Mammals	0

ESA Status	
SC	13
No Status	2
EXP	0
TOTAL SCC	15